

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 September 2003 (25.09.2003)

PCT

(10) International Publication Number  
**WO 03/077918 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 31/4439**,  
C07D 401/12

**Brian, W.** [CA/US]; 126 East Lincoln Avenue, Rahway,  
NJ 07065-0907 (US).

(21) International Application Number: PCT/US03/07074

(74) Common Representative: **MERCK & CO., INC.**; 126  
East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(22) International Filing Date: 7 March 2003 (07.03.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/363,456 12 March 2002 (12.03.2002) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,  
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VC, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **MERCK  
& CO., INC.** [US/US]; 126 East Lincoln Avenue, Rahway,  
NJ 07065-0907 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **SMITH, Nicholas,  
D.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ  
07065-0907 (US). **COSFORD, Nicholas, D. P.** [GB/US];  
126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).  
**REGER, Thomas, R.** [US/US]; 126 East Lincoln Ave-  
nue, Rahway, NJ 07065-0907 (US). **ROPPE, Jeffrey,  
R.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ  
07065-0907 (US). **POON, Steven, F.** [US/US]; 126 East  
Lincoln Avenue, Rahway, NJ 07065-0907 (US). **HUANG,  
Dehua** [CN/US]; 126 East Lincoln Avenue, Rahway, NJ  
07065-0907 (US). **CHEN, Chixu** [US/US]; 126 East Lin-  
coln Avenue, Rahway, NJ 07065-0907 (US). **EASTMAN,**

**Published:**

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

(54) Title: DI-ARYL SUBSTITUTED TETRAZOLE MODULATORS OF METABOTROPIC GLUTAMATE RECEPTOR-5

(57) Abstract: Tetrazole compounds substituted directly, or by a bridge, with i) a heteroaryl moiety containing N adjacent to the point of connection of the heteroaryl and ii) another heteroaryl or aryl ring, with at least one of the rings being further substituted with another ring, are mGluR5 modulators useful in the treatment of psychiatric and mood disorders such as, for example, schizophrenia, anxiety, depression, panic, and bipolar disorder, as well as in the treatment of pain, Parkinson's disease, cognitive dysfunction, epilepsy, circadian rhythm disorders, drug addiction, drug abuse, drug withdrawal, obesity and other diseases.



WO 03/077918 A1

## TITLE OF THE INVENTION

DI-ARYL SUBSTITUTED TETRAZOLE MODULATORS OF METABOTROPIC  
GLUTAMATE RECEPTOR-5

5

## BACKGROUND OF THE INVENTION

## FIELD OF THE INVENTION

The present invention is directed to tetrazole compounds substituted  
10 with i) a heteroaryl ring and ii) another heteroaryl or aryl ring with at least one of the  
rings being further substituted with another ring. In particular, this invention is  
directed to tetrazole compounds substituted directly, or by a bridge, with i) a  
heteroaryl moiety containing N adjacent to the point of connection of the heteroaryl  
and ii) another heteroaryl or aryl ring, with at least one of the rings being further  
15 substituted with another ring, which are metabotropic glutamate receptor – subtype 5  
("mGluR5") modulators useful in the treatment of psychiatric and mood disorders  
such as, for example, schizophrenia, anxiety, depression, panic, bipolar disorder, and  
circadian rhythm disorders, as well as in the treatment of pain, Parkinson's disease,  
cognitive dysfunction, epilepsy, drug addiction, drug abuse, drug withdrawal, obesity  
20 and other diseases.

## RELATED BACKGROUND

A major excitatory neurotransmitter in the mammalian nervous system  
is the glutamate molecule, which binds to neurons, thereby activating cell surface  
25 receptors. Such surface receptors are characterized as either ionotropic or  
metabotropic glutamate receptors. The metabotropic glutamate receptors ("mGluR")  
are G protein-coupled receptors that activate intracellular second messenger systems  
when bound to glutamate. Activation of mGluR results in a variety of cellular  
responses. In particular, mGluR1 and mGluR5 activate phospholipase C, which is  
30 followed by mobilizing intracellular calcium.

Modulation of metabotropic glutamate receptor subtype 5 (mGluR5) is  
useful in the treatment of diseases that affect the nervous system (see for example  
W.P.J.M Spooren et al., *Trends Pharmacol. Sci.*, 22:331-337 (2001) and references  
cited therein). For example, recent evidence demonstrates the involvement of  
35 mGluR5 in nociceptive processes and that modulation of mGluR5 using mGluR5-

selective compounds is useful in the treatment of various pain states, including acute, persistent and chronic pain [K Walker et al., *Neuropharmacology*, 40:1-9 (2001); F. Bordi, A. Ugolini *Brain Res.*, 871:223-233 (2001)], inflammatory pain [K Walker et al., *Neuropharmacology*, 40:10-19 (2001); Bhawe et al. *Nature Neurosci.* 4:417-423 (2001)] and neuropathic pain [Dogrul et al. *Neurosci. Lett.* 292:115-118 (2000)].

Further evidence supports the use of modulators of mGluR5 in the treatment of psychiatric and neurological disorders. For example, mGluR5-selective compounds such as 2-methyl-6-(phenylethynyl)-pyridine ("MPEP") are effective in animal models of mood disorders, including anxiety and depression [W.P.J.M Spoooren et al., *J. Pharmacol. Exp. Ther.*, 295:1267-1275 (2000); E. Tatarczynska et al., *Brit. J. Pharmacol.*, 132:1423-1430 (2001); A. Klodzynska et al, *Pol. J. Pharmacol.*, 132:1423-1430 (2001)]. Gene expression data from humans indicate that modulation of mGluR5 may be useful for the treatment of schizophrenia [T. Ohnuma et al, *Mol. Brain. Res.*, 56:207-217 (1998); *ibid*, *Mol. Brain. Res.*, 85:24-31 (2000)].

Studies have also shown a role for mGluR5, and the potential utility of mGluR5-modulatory compounds, in the treatment of movement disorders such as Parkinson's disease [W.P.J.M Spoooren et al., *Europ. J. Pharmacol.* 406:403-410 (2000); H. Awad et al., *J. Neurosci.* 20:7871-7879 (2000); K. Ossawa et al. *Neuropharmacol.* 41:413-420 (2001)]. Other research supports a role for mGluR5 modulation in the treatment of cognitive dysfunction [G. Riedel et al, *Neuropharmacol.* 39:1943-1951 (2000)], epilepsy [A. Chapman et al, *Neuropharmacol.* 39:1567-1574 (2000)] and neuroprotection [V. Bruno et al, *Neuropharmacol.* 39:2223-2230 (2000)]. Studies with mGluR5 knockout mice and MPEP also suggest that modulation of these receptors may be useful in the treatment of drug addiction, drug abuse and drug withdrawal [C. Chiamulera et al. *Nature Neurosci.* 4:873-874 (2001)].

International Patent Publications WO 01/12627 and WO 99/26927 describe heteropolycyclic compounds and their use as metabotropic glutamate receptor antagonists.

U.S. Patent No. 3,647,809 describes pyridyl-1,2,4-oxadiazole derivatives. U.S. Patent No. 4,022,901 describes 3-pyridyl-5-isothiocyanophenyl oxadiazoles. International Patent Publication WO 98/17652 describes oxadiazoles, WO 97/03967 describes various substituted aromatic compounds, JP 13233767A and WO 94/22846 describe various heterocyclic compounds.

Compounds that include ringed systems are described by various investigators as effective for a variety of therapies and utilities. For example,

International Patent Publication No. WO 98/25883 describes ketobenzamides as calpain inhibitors, European Patent Publication No. EP 811610 and U.S. Patent Nos. 5,679,712, 5,693,672 and 5,747,541 describe substituted benzoylguanidine sodium channel blockers, and U.S. Patent No. 5,736,297 describes ring systems useful as a  
5 photosensitive composition.

However, there remains a need for novel compounds and compositions that therapeutically inhibit mGluR5 with minimal side effects.


#### SUMMARY OF THE INVENTION

10 The present invention is directed to novel tetrazole compounds substituted directly, or by a bridge, with i) a heteroaryl moiety containing N adjacent to the point of connection of the heteroaryl and ii) another heteroaryl or aryl ring, with at least one of the rings being further substituted with another ring, which are  
15 metabotropic glutamate receptor – subtype 5 modulators useful in the treatment of psychiatric and mood disorders such as, for example, schizophrenia, anxiety, depression, bipolar disorders, and panic, as well as in the treatment of pain, Parkinson's disease, cognitive dysfunction, epilepsy, circadian rhythm and sleep disorders – such as shift-work induced sleep disorder and jet-lag, drug addiction, drug  
20 abuse, drug withdrawal, obesity and other diseases. This invention also provides a pharmaceutical composition which includes an effective amount of the novel tetrazole compounds substituted with a heteroaryl moiety, and a pharmaceutically acceptable carrier.

This invention further provides a method of treatment of psychiatric and mood disorders such as, for example, schizophrenia, anxiety, depression, panic,  
25 bipolar disorders, and circadian rhythm and sleep disorders, as well as a method of treatment of pain, Parkinson's disease, cognitive dysfunction, epilepsy, obesity, drug addiction, drug abuse and drug withdrawal by the administration of an effective amount of the novel tetrazole compounds substituted with a heteroaryl moiety.

#### 30 DETAILED DESCRIPTION OF THE INVENTION

The compounds of the present invention are represented by Formula (I):



(I)

or a pharmaceutically acceptable salt thereof, wherein

X and Y each independently is aryl or heteroaryl wherein at least one  
 5 of X and Y is a heteroaryl with N adjacent to the position of attachment to A or B  
 respectively;

X is optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>,  
 -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>,  
 -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>,  
 10 -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or  
 -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are combined to  
 form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1-6</sub>alkyl  
 substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl),  
 15 -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or  
 -N(C<sub>0-6</sub>alkyl)(aryl) groups;

R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl,  
 heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl),  
 20 -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

R<sup>4</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally  
 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl),  
 -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl),  
 -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

25 A is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-,  
 -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

W is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl,  
 30 -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>,

-N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

- Y is optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>,  
 5 -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>,  
 -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>, -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>,  
 -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)R<sup>6</sup>, or  
 -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are combined to  
 form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>alkyl  
 10 substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl),  
 -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or  
 -N(C<sub>0-6</sub>alkyl)(aryl) groups;

- R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl,  
 15 heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>  
 alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;  
 R<sup>8</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally  
 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>  
 20 cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl),  
 -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

- B is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>  
 alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-  
 NR<sup>10</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

- 25 R<sup>9</sup> and R<sup>10</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl,  
 heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>  
 alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

- Z is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>  
 30 alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>  
 alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -  
 N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>,  
 -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -  
 C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

one of W and Z is optionally absent; and  
any N may be an N-oxide.

- In one aspect, the compounds of this invention are represented by
- 5 Formula (I) or a pharmaceutically acceptable salt thereof, wherein
- X is 2-pyridyl optionally substituted with 1-4 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>,  
10 -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or  
15 -N(C<sub>0-6</sub>alkyl)(aryl) groups;
- R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;
- 20 R<sup>4</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;
- A is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;
- 25 W is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>,  
30 -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;
- Y is aryl or heteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>,  
35 -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>,

-NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>,  
 -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are  
 combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1</sub>-  
 6alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 5 substituted with 1-5 independent halogen, -CN, -C<sub>1</sub>-6alkyl, -O(C<sub>0</sub>-6alkyl), -O(C<sub>3</sub>-  
 7cycloalkyl), -O(aryl), -N(C<sub>0</sub>-6alkyl)(C<sub>0</sub>-6alkyl), -N(C<sub>0</sub>-6alkyl)(C<sub>3</sub>-7cycloalkyl), or  
 -N(C<sub>0</sub>-6alkyl)(aryl) groups;

R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is -C<sub>0</sub>-6alkyl, -C<sub>3</sub>-7cycloalkyl,  
 heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 10 halogen, -CN, -C<sub>1</sub>-6alkyl, -O(C<sub>0</sub>-6alkyl), -O(C<sub>3</sub>-7cycloalkyl), -O(aryl), -N(C<sub>0</sub>-  
 6alkyl)(C<sub>0</sub>-6alkyl), -N(C<sub>0</sub>-6alkyl)(C<sub>3</sub>-7cycloalkyl), -N(C<sub>0</sub>-6alkyl)(aryl) substituents;

R<sup>8</sup> is -C<sub>1</sub>-6alkyl, -C<sub>3</sub>-7cycloalkyl, heteroaryl, or aryl; optionally  
 substituted with 1-5 independent halogen, -CN, -C<sub>1</sub>-6alkyl, -O(C<sub>0</sub>-6alkyl), -O(C<sub>3</sub>-  
 7cycloalkyl), -O(aryl), -N(C<sub>0</sub>-6alkyl)(C<sub>0</sub>-6alkyl), -N(C<sub>0</sub>-6alkyl)(C<sub>3</sub>-7cycloalkyl),  
 15 -N(C<sub>0</sub>-6alkyl)(aryl) substituents;

B is -C<sub>0</sub>-4alkyl, -C<sub>0</sub>-2alkyl-SO-C<sub>0</sub>-2alkyl-, -C<sub>0</sub>-2alkyl-SO<sub>2</sub>-C<sub>0</sub>-  
 2alkyl-, -C<sub>0</sub>-2alkyl-CO-C<sub>0</sub>-2alkyl-, -C<sub>0</sub>-2alkyl-NR<sup>10</sup>CO-C<sub>0</sub>-2alkyl-, -C<sub>0</sub>-2alkyl-  
 NR<sup>10</sup>SO<sub>2</sub>-C<sub>0</sub>-2alkyl- or -heteroC<sub>0</sub>-4alkyl;

R<sup>9</sup> and R<sup>10</sup> each independently is -C<sub>0</sub>-6alkyl, -C<sub>3</sub>-7cycloalkyl,  
 20 heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 halogen, -CN, -C<sub>1</sub>-6alkyl, -O(C<sub>0</sub>-6alkyl), -O(C<sub>3</sub>-7cycloalkyl), -O(aryl), -N(C<sub>0</sub>-  
 6alkyl)(C<sub>0</sub>-6alkyl), -N(C<sub>0</sub>-6alkyl)(C<sub>3</sub>-7cycloalkyl), -N(C<sub>0</sub>-6alkyl)(aryl) substituents;

Z is -C<sub>3</sub>-7cycloalkyl, -heteroC<sub>3</sub>-7cycloalkyl, -C<sub>0</sub>-6alkylaryl, or -C<sub>0</sub>-  
 6alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1</sub>-  
 25 6alkyl, -C<sub>1</sub>-6alkenyl, -C<sub>1</sub>-6alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>,  
 -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>,  
 -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or  
 -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

one of W and Z is optionally absent; and  
 30 any N may be an N-oxide.

In an embodiment of this one aspect, the compounds of this invention  
 are represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein



X is 2-pyridyl optionally substituted with 1-4 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1</sub>-6alkyl, -C<sub>1</sub>-6alkenyl, -C<sub>1</sub>-6alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>,  
 5 -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1</sub>-6alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1</sub>-6alkyl, -O(C<sub>0</sub>-6alkyl), -O(C<sub>3</sub>-7cycloalkyl), -O(aryl), -N(C<sub>0</sub>-6alkyl)(C<sub>0</sub>-6alkyl), -N(C<sub>0</sub>-6alkyl)(C<sub>3</sub>-7cycloalkyl), or  
 10 -N(C<sub>0</sub>-6alkyl)(aryl) groups;

R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> each independently is -C<sub>0</sub>-6alkyl, -C<sub>3</sub>-7cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1</sub>-6alkyl, -O(C<sub>0</sub>-6alkyl), -O(C<sub>3</sub>-7cycloalkyl), -O(aryl), -N(C<sub>0</sub>-6alkyl)(C<sub>0</sub>-6alkyl), -N(C<sub>0</sub>-6alkyl)(C<sub>3</sub>-7cycloalkyl), -N(C<sub>0</sub>-6alkyl)(aryl) substituents;  
 15 R<sup>4</sup> is -C<sub>1</sub>-6alkyl, -C<sub>3</sub>-7cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1</sub>-6alkyl, -O(C<sub>0</sub>-6alkyl), -O(C<sub>3</sub>-7cycloalkyl), -O(aryl), -N(C<sub>0</sub>-6alkyl)(C<sub>0</sub>-6alkyl), -N(C<sub>0</sub>-6alkyl)(C<sub>3</sub>-7cycloalkyl), -N(C<sub>0</sub>-6alkyl)(aryl) substituents;

A is -C<sub>0</sub>-4alkyl, -C<sub>0</sub>-2alkyl-SO-C<sub>0</sub>-2alkyl-, -C<sub>0</sub>-2alkyl-SO<sub>2</sub>-C<sub>0</sub>-2alkyl-, -C<sub>0</sub>-2alkyl-CO-C<sub>0</sub>-2alkyl-, -C<sub>0</sub>-2alkyl-NR<sup>9</sup>CO-C<sub>0</sub>-2alkyl-, -C<sub>0</sub>-2alkyl-NR<sup>9</sup>SO<sub>2</sub>-C<sub>0</sub>-2alkyl- or -heteroC<sub>0</sub>-4alkyl;

W is -C<sub>3</sub>-7cycloalkyl, -heteroC<sub>3</sub>-7cycloalkyl, -C<sub>0</sub>-6alkylaryl, or -C<sub>0</sub>-6alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1</sub>-6alkyl, -C<sub>1</sub>-6alkenyl, -C<sub>1</sub>-6alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>,  
 25 -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

Y is phenyl optionally substituted with 1-5 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1</sub>-6alkyl, -C<sub>1</sub>-6alkenyl, -C<sub>1</sub>-6alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>,  
 30 -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>, -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1</sub>-6alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 35 substituted with 1-5 independent halogen, -CN, -C<sub>1</sub>-6alkyl, -O(C<sub>0</sub>-6alkyl), -O(C<sub>3</sub>-

7cycloalkyl), -O(aryl), -N(C<sub>0</sub>-6alkyl)(C<sub>0</sub>-6alkyl), -N(C<sub>0</sub>-6alkyl)(C<sub>3</sub>-7cycloalkyl), or -N(C<sub>0</sub>-6alkyl)(aryl) groups;

R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is -C<sub>0</sub>-6alkyl, -C<sub>3</sub>-7cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 5 halogen, -CN, -C<sub>1</sub>-6alkyl, -O(C<sub>0</sub>-6alkyl), -O(C<sub>3</sub>-7cycloalkyl), -O(aryl), -N(C<sub>0</sub>-6alkyl)(C<sub>0</sub>-6alkyl), -N(C<sub>0</sub>-6alkyl)(C<sub>3</sub>-7cycloalkyl), -N(C<sub>0</sub>-6alkyl)(aryl) substituents;

R<sup>8</sup> is -C<sub>1</sub>-6alkyl, -C<sub>3</sub>-7cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1</sub>-6alkyl, -O(C<sub>0</sub>-6alkyl), -O(C<sub>3</sub>-7cycloalkyl), -O(aryl), -N(C<sub>0</sub>-6alkyl)(C<sub>0</sub>-6alkyl), -N(C<sub>0</sub>-6alkyl)(C<sub>3</sub>-7cycloalkyl),  
 10 -N(C<sub>0</sub>-6alkyl)(aryl) substituents;

B is -C<sub>0</sub>-4alkyl, -C<sub>0</sub>-2alkyl-SO-C<sub>0</sub>-2alkyl-, -C<sub>0</sub>-2alkyl-SO<sub>2</sub>-C<sub>0</sub>-2alkyl-, -C<sub>0</sub>-2alkyl-CO-C<sub>0</sub>-2alkyl-, -C<sub>0</sub>-2alkyl-NR<sup>10</sup>CO-C<sub>0</sub>-2alkyl-, -C<sub>0</sub>-2alkyl-NR<sup>10</sup>SO<sub>2</sub>-C<sub>0</sub>-2alkyl- or -heteroC<sub>0</sub>-4alkyl;

R<sup>9</sup> and R<sup>10</sup> each independently is -C<sub>0</sub>-6alkyl, -C<sub>3</sub>-7cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 15 halogen, -CN, -C<sub>1</sub>-6alkyl, -O(C<sub>0</sub>-6alkyl), -O(C<sub>3</sub>-7cycloalkyl), -O(aryl), -N(C<sub>0</sub>-6alkyl)(C<sub>0</sub>-6alkyl), -N(C<sub>0</sub>-6alkyl)(C<sub>3</sub>-7cycloalkyl), -N(C<sub>0</sub>-6alkyl)(aryl) substituents;

Z is -C<sub>3</sub>-7cycloalkyl, -heteroC<sub>3</sub>-7cycloalkyl, -C<sub>0</sub>-6alkylaryl, or -C<sub>0</sub>-6alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1</sub>-  
 20 6alkyl, -C<sub>1</sub>-6alkenyl, -C<sub>1</sub>-6alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

one of W and Z is optionally absent; and  
 25 any N may be an N-oxide.

In a second aspect, the compounds of this invention are represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

X is aryl or heteroaryl optionally substituted with 1-7 independent  
 30 halogen, -CN, NO<sub>2</sub>, -C<sub>1</sub>-6alkyl, -C<sub>1</sub>-6alkenyl, -C<sub>1</sub>-6alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1</sub>-

6alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or -N(C<sub>0-6</sub>alkyl)(aryl) groups;

5 R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

R<sup>4</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally  
10 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

A is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-  
15 NR<sup>9</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

W is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>,  
-N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>,  
20 -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

Y is 2-pyridyl optionally substituted with 1-4 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>,  
-C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>,  
25 -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or  
30 -N(C<sub>0-6</sub>alkyl)(aryl) groups;

R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent

halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

R<sup>8</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

B is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

R<sup>9</sup> and R<sup>10</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

Z is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

one of W and Z is optionally absent; and  
any N may be an N-oxide.

In a third aspect, the compounds of this invention are represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

X is phenyl optionally substituted with 1-5 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or -N(C<sub>0-6</sub>alkyl)(aryl) groups;

- R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;
- 5 R<sup>4</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;
- 10 A is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;
- W is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>,  
 15 -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;
- Y is aryl or heteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>,  
 20 -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>, -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 25 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or -N(C<sub>0-6</sub>alkyl)(aryl) groups;
- R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 30 halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;
- R<sup>8</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-</sub>

7cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

B is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

R<sup>9</sup> and R<sup>10</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

Z is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

one of W and Z is optionally absent; and  
any N may be an N-oxide.

In an embodiment of this third aspect, the compounds of this invention are represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

X is phenyl optionally substituted with 1-5 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or -N(C<sub>0-6</sub>alkyl)(aryl) groups;

R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

R<sup>4</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

5 A is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

W is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>,  
 10 -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

Y is 2-pyridyl optionally substituted with 1-4 independent halogen,  
 15 -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>, -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 20 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or -N(C<sub>0-6</sub>alkyl)(aryl) groups;

R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 25 halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

R<sup>8</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl),  
 30 -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

B is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

- R<sup>9</sup> and R<sup>10</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;
- 5 Z is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or
- 10 -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;
- one of W and Z is optionally absent; and  
any N may be an N-oxide.

- In a fourth aspect, the compounds of this invention are represented by
- 15 Formula (I) or a pharmaceutically acceptable salt thereof, wherein
- X is aryl or heteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>,
- 20 -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or
- 25 -N(C<sub>0-6</sub>alkyl)(aryl) groups;
- R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;
- 30 R<sup>4</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;



A is  $-C_{0-4}alkyl$ ,  $-C_{0-2}alkyl-SO-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-SO_2-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-CO-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-NR^9CO-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-NR^9SO_2-C_{0-2}alkyl-$  or  $-heteroC_{0-4}alkyl$ ;

W is  $-C_{3-7}cycloalkyl$ ,  $-heteroC_{3-7}cycloalkyl$ ,  $-C_{0-6}alkylaryl$ , or  $-C_{0-6}alkylheteroaryl$  optionally substituted with 1-7 independent halogen,  $-CN$ ,  $NO_2$ ,  $-C_{1-6}alkyl$ ,  $-C_{1-6}alkenyl$ ,  $-C_{1-6}alkynyl$ ,  $-OR^1$ ,  $-NR^1R^2$ ,  $-C(=NR^1)NR^2R^3$ ,  $-N(=NR^1)NR^2R^3$ ,  $-NR^1COR^2$ ,  $-NR^1CO_2R^2$ ,  $-NR^1SO_2R^4$ ,  $-NR^1CONR^2R^3$ ,  $-SR^4$ ,  $-SOR^4$ ,  $-SO_2R^4$ ,  $-SO_2NR^1R^2$ ,  $-COR^1$ ,  $-CO_2R^1$ ,  $-CONR^1R^2$ ,  $-C(=NR^1)R^2$ , or  $-C(=NOR^1)R^2$  substituents;

Y is 1,3-thiazolyl optionally substituted with 1-2 independent halogen,  $-CN$ ,  $NO_2$ ,  $-C_{1-6}alkyl$ ,  $-C_{1-6}alkenyl$ ,  $-C_{1-6}alkynyl$ ,  $-OR^5$ ,  $-NR^5R^6$ ,  $-C(=NR^5)NR^6R^7$ ,  $-N(=NR^5)NR^6R^7$ ,  $-NR^5COR^6$ ,  $-NR^5CO_2R^6$ ,  $-NR^5SO_2R^8$ ,  $-NR^5CONR^6R^7$ ,  $-SR^8$ ,  $-SOR^8$ ,  $-SO_2R^8$ ,  $-SO_2NR^5R^6$ ,  $-COR^5$ ,  $-CO_2R^5$ ,  $-CONR^5R^6$ ,  $-C(=NR^5)R^6$ , or  $-C(=NOR^5)R^6$  substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the  $-C_{1-6}alkyl$  substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen,  $-CN$ ,  $-C_{1-6}alkyl$ ,  $-O(C_{0-6}alkyl)$ ,  $-O(C_{3-7}cycloalkyl)$ ,  $-O(aryl)$ ,  $-N(C_{0-6}alkyl)(C_{0-6}alkyl)$ ,  $-N(C_{0-6}alkyl)(C_{3-7}cycloalkyl)$ , or  $-N(C_{0-6}alkyl)(aryl)$  groups;

$R^5$ ,  $R^6$ , and  $R^7$  each independently is  $-C_{0-6}alkyl$ ,  $-C_{3-7}cycloalkyl$ , heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen,  $-CN$ ,  $-C_{1-6}alkyl$ ,  $-O(C_{0-6}alkyl)$ ,  $-O(C_{3-7}cycloalkyl)$ ,  $-O(aryl)$ ,  $-N(C_{0-6}alkyl)(C_{0-6}alkyl)$ ,  $-N(C_{0-6}alkyl)(C_{3-7}cycloalkyl)$ ,  $-N(C_{0-6}alkyl)(aryl)$  substituents;

$R^8$  is  $-C_{1-6}alkyl$ ,  $-C_{3-7}cycloalkyl$ , heteroaryl, or aryl; optionally substituted with 1-5 independent halogen,  $-CN$ ,  $-C_{1-6}alkyl$ ,  $-O(C_{0-6}alkyl)$ ,  $-O(C_{3-7}cycloalkyl)$ ,  $-O(aryl)$ ,  $-N(C_{0-6}alkyl)(C_{0-6}alkyl)$ ,  $-N(C_{0-6}alkyl)(C_{3-7}cycloalkyl)$ ,  $-N(C_{0-6}alkyl)(aryl)$  substituents;

B is  $-C_{0-4}alkyl$ ,  $-C_{0-2}alkyl-SO-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-SO_2-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-CO-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-NR^{10}CO-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-NR^{10}SO_2-C_{0-2}alkyl-$  or  $-heteroC_{0-4}alkyl$ ;

$R^9$  and  $R^{10}$  each independently is  $-C_{0-6}alkyl$ ,  $-C_{3-7}cycloalkyl$ , heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen,  $-CN$ ,  $-C_{1-6}alkyl$ ,  $-O(C_{0-6}alkyl)$ ,  $-O(C_{3-7}cycloalkyl)$ ,  $-O(aryl)$ ,  $-N(C_{0-6}alkyl)(C_{0-6}alkyl)$ ,  $-N(C_{0-6}alkyl)(C_{3-7}cycloalkyl)$ ,  $-N(C_{0-6}alkyl)(aryl)$  substituents;

Z is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>,  
 -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>,  
 5 -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or  
 -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

one of W and Z is optionally absent; and

any N may be an N-oxide.

10 In an embodiment of this fourth aspect, the compounds of this invention are represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

X is phenyl optionally substituted with 1-5 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>,  
 15 -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>,  
 -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or  
 -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1-6</sub>alkyl  
 substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 20 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or  
 -N(C<sub>0-6</sub>alkyl)(aryl) groups;

R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 25 halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

R<sup>4</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl),  
 30 -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

A is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

W is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>,  
 -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>,  
 5 -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or  
 -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

Y is 1,3-thiazolyl optionally substituted with 1-2 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>,  
 -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>,  
 10 -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>,  
 -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are  
 combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>  
 6alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>  
 15 7cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or  
 -N(C<sub>0-6</sub>alkyl)(aryl) groups;

R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>  
 20 6alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

R<sup>8</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>  
 7cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl),  
 -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

B is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-  
 25 NR<sup>10</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

R<sup>9</sup> and R<sup>10</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>  
 30 6alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

Z is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>  
 6alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>,

-N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>,  
 -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or  
 -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

one of W and Z is optionally absent; and

5 any N may be an N-oxide.

In a fifth aspect, the compounds of this invention are represented by  
 Formula (I) or a pharmaceutically acceptable salt thereof, wherein

X is aryl or heteroaryl optionally substituted with 1-7 independent  
 10 halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>,  
 -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>,  
 -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>,  
 -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are  
 15 combined to form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1-6</sub>  
 6alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>  
 cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or  
 -N(C<sub>0-6</sub>alkyl)(aryl) groups;

R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl,  
 20 heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>  
 6alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

R<sup>4</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally  
 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>  
 25 7cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl),  
 -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

A is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>  
 2alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-  
 NR<sup>9</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

W is -C<sub>0-6</sub>alkylaryl or -C<sub>0-6</sub>alkylheteroaryl optionally substituted  
 30 with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl,  
 -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>,  
 -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>,  
 -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

Y is aryl or heteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>, -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>,  
 5 -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or  
 10 -N(C<sub>0-6</sub>alkyl)(aryl) groups;

R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;  
 15 R<sup>8</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

B is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;  
 20

R<sup>9</sup> and R<sup>10</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;  
 25

Z is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>,  
 30 -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

one of W and Z is optionally absent; and  
 any N may be an N-oxide.

In a sixth aspect, the compounds of this invention are represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

X is aryl or heteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>,  
 5 -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>,  
 -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>,  
 -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are  
 combined to form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1-6</sub>  
 10 alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl),  
 -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or  
 -N(C<sub>0-6</sub>alkyl)(aryl) groups;

R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl,  
 heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 15 halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>  
 alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

R<sup>4</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally  
 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>  
 7cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl),  
 20 -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

A is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>  
 alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-  
 NR<sup>9</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

W is -C<sub>0-6</sub>alkylaryl or -C<sub>0-6</sub>alkylheteroaryl optionally substituted  
 25 with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl,  
 -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>,  
 -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>,  
 -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

Y is pyrazolyl optionally substituted with 1-3 independent halogen,  
 30 -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>,  
 -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>,  
 -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>,  
 -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are  
 combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>  
 35 alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further

substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or -N(C<sub>0-6</sub>alkyl)(aryl) groups;

R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

R<sup>8</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

B is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

R<sup>9</sup> and R<sup>10</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

Z is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

one of W and Z is optionally absent; and  
any N may be an N-oxide.

In an embodiment of this sixth aspect, the compounds of this invention are represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

X is phenyl optionally substituted with 1-5 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are combined to form

a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or -N(C<sub>0-6</sub>alkyl)(aryl) groups;

R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

R<sup>4</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

A is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

W is -C<sub>0-6</sub>alkylaryl or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

Y is pyrazolyl optionally substituted with 1-3 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>, -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or -N(C<sub>0-6</sub>alkyl)(aryl) groups;

R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent



- halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;
- R<sup>8</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;
- B is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;
- R<sup>9</sup> and R<sup>10</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;
- Z is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;
- one of W and Z is optionally absent; and  
any N may be an N-oxide.

- In a seventh aspect, the compounds of this invention are represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein
- X is aryl or heteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or -N(C<sub>0-6</sub>alkyl)(aryl) groups;

- R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;
- 5 R<sup>4</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;
- 10 A is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;
- 15 W is -C<sub>0-6</sub>alkylaryl or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;
- 20 Y is imidazolyl optionally substituted with 1-3 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>, -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or -N(C<sub>0-6</sub>alkyl)(aryl) groups;
- 25 R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;
- 30 R<sup>8</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

B is  $-C_{0-4}alkyl$ ,  $-C_{0-2}alkyl-SO-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-SO_2-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-CO-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-NR^{10}CO-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-NR^{10}SO_2-C_{0-2}alkyl-$  or  $-heteroC_{0-4}alkyl$ ;

$R^9$  and  $R^{10}$  each independently is  $-C_{0-6}alkyl$ ,  $-C_{3-7}cycloalkyl$ ,  
 5 heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen,  $-CN$ ,  $-C_{1-6}alkyl$ ,  $-O(C_{0-6}alkyl)$ ,  $-O(C_{3-7}cycloalkyl)$ ,  $-O(aryl)$ ,  $-N(C_{0-6}alkyl)(C_{0-6}alkyl)$ ,  $-N(C_{0-6}alkyl)(C_{3-7}cycloalkyl)$ ,  $-N(C_{0-6}alkyl)(aryl)$  substituents;

Z is  $-C_{3-7}cycloalkyl$ ,  $-heteroC_{3-7}cycloalkyl$ ,  $-C_{0-6}alkylaryl$ , or  $-C_{0-6}alkylheteroaryl$  optionally substituted with 1-7 independent halogen,  $-CN$ ,  $NO_2$ ,  $-C_{1-6}alkyl$ ,  $-C_{1-6}alkenyl$ ,  $-C_{1-6}alkynyl$ ,  $-OR^1$ ,  $-NR^1R^2$ ,  $-C(=NR^1)NR^2R^3$ ,  
 10  $-N(=NR^1)NR^2R^3$ ,  $-NR^1COR^2$ ,  $-NR^1CO_2R^2$ ,  $-NR^1SO_2R^4$ ,  $-NR^1CONR^2R^3$ ,  $-SR^4$ ,  $-SOR^4$ ,  $-SO_2R^4$ ,  $-SO_2NR^1R^2$ ,  $-COR^1$ ,  $-CO_2R^1$ ,  $-CONR^1R^2$ ,  $-C(=NR^1)R^2$ , or  $-C(=NOR^1)R^2$  substituents;

one of W and Z is optionally absent; and  
 15 any N may be an N-oxide.

In an embodiment of this seventh aspect, the compounds of this invention are represented by Formula (I) or a pharmaceutically acceptable salt thereof, wherein

20 X is phenyl optionally substituted with 1-5 independent halogen,  $-CN$ ,  $NO_2$ ,  $-C_{1-6}alkyl$ ,  $-C_{1-6}alkenyl$ ,  $-C_{1-6}alkynyl$ ,  $-OR^1$ ,  $-NR^1R^2$ ,  $-C(=NR^1)NR^2R^3$ ,  $-N(=NR^1)NR^2R^3$ ,  $-NR^1COR^2$ ,  $-NR^1CO_2R^2$ ,  $-NR^1SO_2R^4$ ,  $-NR^1CONR^2R^3$ ,  $-SR^4$ ,  $-SOR^4$ ,  $-SO_2R^4$ ,  $-SO_2NR^1R^2$ ,  $-COR^1$ ,  $-CO_2R^1$ ,  $-CONR^1R^2$ ,  $-C(=NR^1)R^2$ , or  $-C(=NOR^1)R^2$  substituents, wherein optionally two substituents are combined to  
 25 form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the  $-C_{1-6}alkyl$  substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen,  $-CN$ ,  $-C_{1-6}alkyl$ ,  $-O(C_{0-6}alkyl)$ ,  $-O(C_{3-7}cycloalkyl)$ ,  $-O(aryl)$ ,  $-N(C_{0-6}alkyl)(C_{0-6}alkyl)$ ,  $-N(C_{0-6}alkyl)(C_{3-7}cycloalkyl)$ , or  $-N(C_{0-6}alkyl)(aryl)$  groups;

30  $R^1$ ,  $R^2$ , and  $R^3$  each independently is  $-C_{0-6}alkyl$ ,  $-C_{3-7}cycloalkyl$ , heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen,  $-CN$ ,  $-C_{1-6}alkyl$ ,  $-O(C_{0-6}alkyl)$ ,  $-O(C_{3-7}cycloalkyl)$ ,  $-O(aryl)$ ,  $-N(C_{0-6}alkyl)(C_{0-6}alkyl)$ ,  $-N(C_{0-6}alkyl)(C_{3-7}cycloalkyl)$ ,  $-N(C_{0-6}alkyl)(aryl)$  substituents;

R<sup>4</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

5 A is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>9</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

10 W is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

15 Y is imidazolyl optionally substituted with 1-3 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>, -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or -N(C<sub>0-6</sub>alkyl)(aryl) groups;

20 R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

25 R<sup>8</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

30 B is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

- R<sup>9</sup> and R<sup>10</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;
- 5 Z is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -
- 10 C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;
- one of W and Z is optionally absent; and  
any N may be an N-oxide.

- In an eighth aspect, the compounds of this invention are represented by
- 15 Formula (I) or a pharmaceutically acceptable salt thereof, wherein

- X is 3-pyridyl optionally substituted with 1-4 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>,
- 20 -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or
- 25 -N(C<sub>0-6</sub>alkyl)(aryl) groups;

- R<sup>1</sup>, R<sup>2</sup>, and R<sup>3</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;
- 30 R<sup>4</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

A is  $-C_{0-4}alkyl$ ,  $-C_{0-2}alkyl-SO-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-SO_2-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-CO-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-NR^9CO-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-NR^9SO_2-C_{0-2}alkyl-$  or  $-heteroC_{0-4}alkyl$ ;

W is  $-C_{3-7}cycloalkyl$ ,  $-heteroC_{3-7}cycloalkyl$ ,  $-C_{0-6}alkylaryl$ , or  $-C_{0-6}alkylheteroaryl$  optionally substituted with 1-7 independent halogen,  $-CN$ ,  $NO_2$ ,  $-C_{1-6}alkyl$ ,  $-C_{1-6}alkenyl$ ,  $-C_{1-6}alkynyl$ ,  $-OR^1$ ,  $-NR^1R^2$ ,  $-C(=NR^1)NR^2R^3$ ,  $-N(=NR^1)NR^2R^3$ ,  $-NR^1COR^2$ ,  $-NR^1CO_2R^2$ ,  $-NR^1SO_2R^4$ ,  $-NR^1CONR^2R^3$ ,  $-SR^4$ ,  $-SOR^4$ ,  $-SO_2R^4$ ,  $-SO_2NR^1R^2$ ,  $-COR^1$ ,  $-CO_2R^1$ ,  $-CONR^1R^2$ ,  $-C(=NR^1)R^2$ , or  $-C(=NOR^1)R^2$  substituents;

Y is aryl or heteroaryl optionally substituted with 1-7 independent halogen,  $-CN$ ,  $NO_2$ ,  $-C_{1-6}alkyl$ ,  $-C_{1-6}alkenyl$ ,  $-C_{1-6}alkynyl$ ,  $-OR^5$ ,  $-NR^5R^6$ ,  $-C(=NR^5)NR^6R^7$ ,  $-N(=NR^5)NR^6R^7$ ,  $-NR^5COR^6$ ,  $-NR^5CO_2R^6$ ,  $-NR^5SO_2R^8$ ,  $-NR^5CONR^6R^7$ ,  $-SR^8$ ,  $-SOR^8$ ,  $-SO_2R^8$ ,  $-SO_2NR^5R^6$ ,  $-COR^5$ ,  $-CO_2R^5$ ,  $-CONR^5R^6$ ,  $-C(=NR^5)R^6$ , or  $-C(=NOR^5)R^6$  substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the  $-C_{1-6}alkyl$  substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen,  $-CN$ ,  $-C_{1-6}alkyl$ ,  $-O(C_{0-6}alkyl)$ ,  $-O(C_{3-7}cycloalkyl)$ ,  $-O(aryl)$ ,  $-N(C_{0-6}alkyl)(C_{0-6}alkyl)$ ,  $-N(C_{0-6}alkyl)(C_{3-7}cycloalkyl)$ , or  $-N(C_{0-6}alkyl)(aryl)$  groups;

$R^5$ ,  $R^6$ , and  $R^7$  each independently is  $-C_{0-6}alkyl$ ,  $-C_{3-7}cycloalkyl$ , heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen,  $-CN$ ,  $-C_{1-6}alkyl$ ,  $-O(C_{0-6}alkyl)$ ,  $-O(C_{3-7}cycloalkyl)$ ,  $-O(aryl)$ ,  $-N(C_{0-6}alkyl)(C_{0-6}alkyl)$ ,  $-N(C_{0-6}alkyl)(C_{3-7}cycloalkyl)$ ,  $-N(C_{0-6}alkyl)(aryl)$  substituents;

$R^8$  is  $-C_{1-6}alkyl$ ,  $-C_{3-7}cycloalkyl$ , heteroaryl, or aryl; optionally substituted with 1-5 independent halogen,  $-CN$ ,  $-C_{1-6}alkyl$ ,  $-O(C_{0-6}alkyl)$ ,  $-O(C_{3-7}cycloalkyl)$ ,  $-O(aryl)$ ,  $-N(C_{0-6}alkyl)(C_{0-6}alkyl)$ ,  $-N(C_{0-6}alkyl)(C_{3-7}cycloalkyl)$ ,  $-N(C_{0-6}alkyl)(aryl)$  substituents;

B is  $-C_{0-4}alkyl$ ,  $-C_{0-2}alkyl-SO-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-SO_2-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-CO-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-NR^{10}CO-C_{0-2}alkyl-$ ,  $-C_{0-2}alkyl-NR^{10}SO_2-C_{0-2}alkyl-$  or  $-heteroC_{0-4}alkyl$ ;

$R^9$  and  $R^{10}$  each independently is  $-C_{0-6}alkyl$ ,  $-C_{3-7}cycloalkyl$ , heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen,  $-CN$ ,  $-C_{1-6}alkyl$ ,  $-O(C_{0-6}alkyl)$ ,  $-O(C_{3-7}cycloalkyl)$ ,  $-O(aryl)$ ,  $-N(C_{0-6}alkyl)(C_{0-6}alkyl)$ ,  $-N(C_{0-6}alkyl)(C_{3-7}cycloalkyl)$ ,  $-N(C_{0-6}alkyl)(aryl)$  substituents;

Z is  $-C_{3-7}$ cycloalkyl,  $-\text{hetero}C_{3-7}$ cycloalkyl,  $-C_{0-6}$ alkylaryl, or  $-C_{0-6}$ alkylheteroaryl optionally substituted with 1-7 independent halogen,  $-\text{CN}$ ,  $\text{NO}_2$ ,  $-C_{1-6}$ alkyl,  $-C_{1-6}$ alkenyl,  $-C_{1-6}$ alkynyl,  $-\text{OR}^1$ ,  $-\text{NR}^1\text{R}^2$ ,  $-\text{C}(=\text{NR}^1)\text{NR}^2\text{R}^3$ ,  
 5  $-\text{N}(=\text{NR}^1)\text{NR}^2\text{R}^3$ ,  $-\text{NR}^1\text{COR}^2$ ,  $-\text{NR}^1\text{CO}_2\text{R}^2$ ,  $-\text{NR}^1\text{SO}_2\text{R}^4$ ,  $-\text{NR}^1\text{CONR}^2\text{R}^3$ ,  $-\text{SR}^4$ ,  
 $-\text{SOR}^4$ ,  $-\text{SO}_2\text{R}^4$ ,  $-\text{SO}_2\text{NR}^1\text{R}^2$ ,  $-\text{COR}^1$ ,  $-\text{CO}_2\text{R}^1$ ,  $-\text{CONR}^1\text{R}^2$ ,  $-\text{C}(=\text{NR}^1)\text{R}^2$ , or  
 $-\text{C}(=\text{NOR}^1)\text{R}^2$  substituents;

one of W and Z is optionally absent; and

any N may be an N-oxide.

As used herein, "alkyl" as well as other groups having the prefix "alk" such as, for example, alkoxy, alkanoyl, alkenyl, alkynyl and the like, means carbon chains which may be linear or branched or combinations thereof. Examples of alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, sec- and tert-butyl, pentyl, hexyl, heptyl and the like. "Alkenyl", "alkynyl" and other like terms include carbon chains containing at least one unsaturated C-C bond.

15 The term "cycloalkyl" means carbocycles containing no heteroatoms, and includes mono-, bi- and tricyclic saturated carbocycles, as well as fused ring systems. Such fused ring systems can include one ring that is partially or fully unsaturated such as a benzene ring to form fused ring systems such as benzofused carbocycles. Cycloalkyl includes such fused ring systems as spirofused ring systems.  
 20 Examples of cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, decahydronaphthalene, adamantane, indanyl, indenyl, fluorenyl, 1,2,3,4-tetrahydronaphthalene and the like. Similarly, "cycloalkenyl" means carbocycles containing no heteroatoms and at least one non-aromatic C-C double bond, and include mono-, bi- and tricyclic partially saturated carbocycles, as well as benzofused  
 25 cycloalkenes. Examples of cycloalkenyl include cyclohexenyl, indenyl, and the like.

The term "aryl" means an aromatic substituent which is a single ring or multiple rings fused together. When formed of multiple rings, at least one of the constituent rings is aromatic. The preferred aryl substituents are phenyl and naphthyl groups.

30 The term "cycloalkyloxy" unless specifically stated otherwise includes a cycloalkyl group connected by a short  $C_{1-2}$ alkyl length to the oxy connecting atom.

The term " $C_{0-6}$ alkyl" includes alkyls containing 6, 5, 4, 3, 2, 1, or no carbon atoms. An alkyl with no carbon atoms is a hydrogen atom substituent when the alkyl is a terminal group and is a direct bond when the alkyl is a bridging group.

The term "hetero" unless specifically stated otherwise includes one or more O, S, or N atoms. For example, heterocycloalkyl and heteroaryl include ring systems that contain one or more O, S, or N atoms in the ring, including mixtures of such atoms. The hetero atoms replace ring carbon atoms. Thus, for example, a heterocycloC<sub>5</sub>alkyl is a five-member ring containing from 4 to no carbon atoms. Examples of heteroaryls include pyridinyl, quinolinyl, isoquinolinyl, pyridazinyl, pyrimidinyl, pyrazinyl, quinoxalinyl, furyl, benzofuryl, dibenzofuryl, thienyl, benzthienyl, pyrrolyl, indolyl, pyrazolyl, indazolyl, oxazolyl, benzoxazolyl, isoxazolyl, thiazolyl, benzothiazolyl, isothiazolyl, imidazolyl, benzimidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, and tetrazolyl. Examples of heterocycloalkyls include azetidiny, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, tetrahydrofuranyl, imidazolinyl, pyrrolidin-2-one, piperidin-2-one, and thiomorpholinyl.

The term "heteroC<sub>0-4</sub>alkyl" means a heteroalkyl containing 3, 2, 1, or no carbon atoms. However, at least one heteroatom must be present. Thus, as an example, a heteroC<sub>0-4</sub>alkyl having no carbon atoms but one N atom would be a -NH- if a bridging group and a -NH<sub>2</sub> if a terminal group. Analogous bridging or terminal groups are clear for an O or S heteroatom.

The term "amine" unless specifically stated otherwise includes primary, secondary and tertiary amines substituted with C<sub>0-6</sub>alkyl.

The term "carbonyl" unless specifically stated otherwise includes a C<sub>0-6</sub>alkyl substituent group when the carbonyl is terminal.

The term "halogen" includes fluorine, chlorine, bromine and iodine atoms.

The term "optionally substituted" is intended to include both substituted and unsubstituted. Thus, for example, optionally substituted aryl could represent a pentafluorophenyl or a phenyl ring. Further, optionally substituted multiple moieties such as, for example, alkylaryl are intended to mean that the aryl and the aryl groups are optionally substituted. If only one of the multiple moieties is optionally substituted then it will be specifically recited such as "an alkylaryl, the aryl optionally substituted with halogen or hydroxyl."

Compounds described herein contain one or more double bonds and may thus give rise to cis/trans isomers as well as other conformational isomers. The present invention includes all such possible isomers as well as mixtures of such isomers.



Compounds described herein can contain one or more asymmetric centers and may thus give rise to diastereomers and optical isomers. The present invention includes all such possible diastereomers as well as their racemic mixtures, their substantially pure resolved enantiomers, all possible geometric isomers, and pharmaceutically acceptable salts thereof. The above Formula I is shown without a definitive stereochemistry at certain positions. The present invention includes all stereoisomers of Formula I and pharmaceutically acceptable salts thereof. Further, mixtures of stereoisomers as well as isolated specific stereoisomers are also included. During the course of the synthetic procedures used to prepare such compounds, or in using racemization or epimerization procedures known to those skilled in the art, the products of such procedures can be a mixture of stereoisomers.

The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic bases or acids. When the compound of the present invention is acidic, its corresponding salt can be conveniently prepared from pharmaceutically acceptable non-toxic bases, including inorganic bases and organic bases. Salts derived from such inorganic bases include aluminum, ammonium, calcium, copper (ic and ous), ferric, ferrous, lithium, magnesium, manganese (ic and ous), potassium, sodium, zinc and the like salts. Particularly preferred are the ammonium, calcium, magnesium, potassium and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, as well as cyclic amines and substituted amines such as naturally occurring and synthesized substituted amines. Other pharmaceutically acceptable organic non-toxic bases from which salts can be formed include ion exchange resins such as, for example, arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine and the like.

When the compound of the present invention is basic, its corresponding salt can be conveniently prepared from pharmaceutically acceptable non-toxic acids, including inorganic and organic acids. Such acids include, for example, acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethanesulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic,

malic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic acid and the like. Particularly preferred are citric, hydrobromic, hydrochloric, maleic, phosphoric, sulfuric, and tartaric acids.

The pharmaceutical compositions of the present invention comprise a  
5 compound represented by Formula I (or pharmaceutically acceptable salts thereof) as  
an active ingredient, a pharmaceutically acceptable carrier and optionally other  
therapeutic ingredients or adjuvants. Such additional therapeutic ingredients include,  
for example, i) opiate agonists or antagonists, ii) calcium channel antagonists, iii)  
5HT receptor agonists or antagonists iv) sodium channel antagonists, v) NMDA  
10 receptor agonists or antagonists, vi) COX-2 selective inhibitors, vii) NK1 antagonists,  
viii) non-steroidal anti-inflammatory drugs ("NSAID"), ix) GABA-A receptor  
modulators, x) dopamine agonists or antagonists, xi) selective serotonin reuptake  
inhibitors ("SSRI") and/or selective serotonin and norepinephrine reuptake inhibitors  
("SSNRI"), xii) tricyclic antidepressant drugs, xiv) norepinephrine modulators, xv) L-  
15 DOPA, xvi) buspirone, xvii) lithium, xviii) valproate, ix) neurontin (gabapentin), xx)  
olanzapine, xxi) nicotinic agonists or antagonists including nicotine, xxii) muscarinic  
agonists or antagonists, xxiii) heroin substituting drugs such as methadone, levo-  
alpha-acetylmethadol, buprenorphine and naltrexone, and xxiv) disulfiram and  
acamprosate. The compositions include compositions suitable for oral, rectal, topical,  
20 and parenteral (including subcutaneous, intramuscular, and intravenous)  
administration, although the most suitable route in any given case will depend on the  
particular host, and nature and severity of the conditions for which the active  
ingredient is being administered. The pharmaceutical compositions may be  
conveniently presented in unit dosage form and prepared by any of the methods well  
25 known in the art of pharmacy.

Creams, ointments, jellies, solutions, or suspensions containing the  
compound of Formula I can be employed for topical use. Mouth washes and gargles  
are included within the scope of topical use for the purposes of this invention.

Dosage levels from about 0.01mg/kg to about 140mg/kg of body  
30 weight per day are useful in the treatment of psychiatric and mood disorders such as,  
for example, schizophrenia, anxiety, depression, panic, bipolar disorders, and  
circadian disorders, as well as being useful in the treatment of pain which are  
responsive to mGluR5 inhibition, or alternatively about 0.5mg to about 7g per patient  
per day. For example, schizophrenia, anxiety, depression, and panic may be  
35 effectively treated by the administration of from about 0.01mg to 75mg of the

compound per kilogram of body weight per day, or alternatively about 0.5mg to about 3.5g per patient per day. Pain may be effectively treated by the administration of from about 0.01mg to 125mg of the compound per kilogram of body weight per day, or alternatively about 0.5mg to about 5.5g per patient per day. Further, it is understood  
5 that the mGluR5 inhibiting compounds of this invention can be administered at prophylactically effective dosage levels to prevent the above-recited conditions.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a formulation intended for  
10 the oral administration to humans may conveniently contain from about 0.5mg to about 5g of active agent, compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95 percent of the total composition. Unit dosage forms will generally contain between from about 1mg to about 1000mg of the active ingredient, typically 25mg, 50mg, 100mg, 200mg, 300mg,  
15 400mg, 500mg, 600mg, 800mg or 1000mg.

It is understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

20 In practice, the compounds represented by Formula I, or pharmaceutically acceptable salts thereof, of this invention can be combined as the active ingredient in intimate admixture with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, e.g.,  
25 oral or parenteral (including intravenous). Thus, the pharmaceutical compositions of the present invention can be presented as discrete units suitable for oral administration such as capsules, cachets or tablets each containing a predetermined amount of the active ingredient. Further, the compositions can be presented as a powder, as granules, as a solution, as a suspension in an aqueous liquid, as a non-aqueous liquid,  
30 as an oil-in-water emulsion or as a water-in-oil liquid emulsion. In addition to the common dosage forms set out above, the compound represented by Formula I, or pharmaceutically acceptable salts thereof, may also be administered by controlled release means and/or delivery devices. The compositions may be prepared by any of the methods of pharmacy. In general, such methods include a step of bringing into  
35 association the active ingredient with the carrier that constitutes one or more

necessary ingredients. In general, the compositions are prepared by uniformly and intimately admixing the active ingredient with liquid carriers or finely divided solid carriers or both. The product can then be conveniently shaped into the desired presentation.

5                   Thus, the pharmaceutical compositions of this invention may include a pharmaceutically acceptable carrier and a compound or a pharmaceutically acceptable salt of Formula I. The compounds of Formula I, or pharmaceutically acceptable salts thereof, can also be included in pharmaceutical compositions in combination with one or more other therapeutically active compounds.

10                   The pharmaceutical carrier employed can be, for example, a solid, liquid, or gas. Examples of solid carriers include lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, and stearic acid. Examples of liquid carriers are sugar syrup, peanut oil, olive oil, and water. Examples of gaseous carriers include carbon dioxide and nitrogen.

15                   In preparing the compositions for oral dosage form, any convenient pharmaceutical media may be employed. For example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like may be used to form oral liquid preparations such as suspensions, elixirs and solutions; while carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants,  
20                   binders, disintegrating agents, and the like may be used to form oral solid preparations such as powders, capsules and tablets. Because of their ease of administration, tablets and capsules are the preferred oral dosage units whereby solid pharmaceutical carriers are employed. Optionally, tablets may be coated by standard aqueous or nonaqueous techniques

25                   A tablet containing the composition of this invention may be prepared by compression or molding, optionally with one or more accessory ingredients or adjuvants. Compressed tablets may be prepared by compressing, in a suitable machine, the active ingredient in a free-flowing form such as powder or granules, optionally mixed with a binder, lubricant, inert diluent, surface active or dispersing  
30                   agent. Molded tablets may be made by molding in a suitable machine, a mixture of the powdered compound moistened with an inert liquid diluent. Each tablet preferably contains from about 0.1mg to about 500mg of the active ingredient and each cachet or capsule preferably containing from about 0.1mg to about 500mg of the active ingredient. Thus, a tablet, cachet, or capsule conveniently contains 0.1mg,  
35                   1mg, 5mg, 25mg, 50mg, 100mg, 200mg, 300mg, 400mg, or 500mg of the active

ingredient taken one or two tablets, cachets, or capsules, once, twice, or three times daily.

Pharmaceutical compositions of the present invention suitable for parenteral administration may be prepared as solutions or suspensions of the active compounds in water. A suitable surfactant can be included such as, for example, hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof in oils. Further, a preservative can be included to prevent the detrimental growth of microorganisms.

Pharmaceutical compositions of the present invention suitable for injectable use include sterile aqueous solutions or dispersions. Furthermore, the compositions can be in the form of sterile powders for the extemporaneous preparation of such sterile injectable solutions or dispersions. In all cases, the final injectable form must be sterile and must be effectively fluid for easy syringability. The pharmaceutical compositions must be stable under the conditions of manufacture and storage; thus, preferably should be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g. glycerol, propylene glycol and liquid polyethylene glycol), vegetable oils, and suitable mixtures thereof.

Pharmaceutical compositions of the present invention can be in a form suitable for topical use such as, for example, an aerosol, cream, ointment, lotion, dusting powder, or the like. Further, the compositions can be in a form suitable for use in transdermal devices. These formulations may be prepared, utilizing a compound represented by Formula I of this invention, or pharmaceutically acceptable salts thereof, via conventional processing methods. As an example, a cream or ointment is prepared by mixing hydrophilic material and water, together with about 5 wt% to about 10 wt% of the compound, to produce a cream or ointment having a desired consistency.

Pharmaceutical compositions of this invention can be in a form suitable for rectal administration wherein the carrier is a solid. It is preferable that the mixture forms unit dose suppositories. Suitable carriers include cocoa butter and other materials commonly used in the art. The suppositories may be conveniently formed by first admixing the composition with the softened or melted carrier(s) followed by chilling and shaping in moulds.

In addition to the aforementioned carrier ingredients, the pharmaceutical formulations described above may include, as appropriate, one or

more additional carrier ingredients such as diluents, buffers, flavoring agents, binders, surface-active agents, thickeners, lubricants, preservatives (including anti-oxidants) and the like. Furthermore, other adjuvants can be included to render the formulation isotonic with the blood of the intended recipient. Compositions containing a  
5 compound described by Formula I, or pharmaceutically acceptable salts thereof, may also be prepared in powder or liquid concentrate form.

The compounds and pharmaceutical compositions of this invention have been found to exhibit biological activity as mGluR5 inhibitors. Accordingly, another aspect of the invention is the treatment in mammals of, for example,  
10 schizophrenia, anxiety, depression, panic, bipolar disorders, circadian rhythm and sleep disorders, pain, Parkinson's disease, cognitive dysfunction, epilepsy, obesity, drug addiction, drug abuse and drug withdrawal – maladies that are amenable to amelioration through inhibition of mGluR5 – by the administration of an effective amount of the compounds of this invention. The term “mammals” includes humans,  
15 as well as other animals such as, for example, dogs, cats, horses, pigs, and cattle. Accordingly, it is understood that the treatment of mammals other than humans is the treatment of clinical correlating afflictions to those above recited examples that are human afflictions.

Further, as described above, the compound of this invention can be  
20 utilized in combination with other therapeutic compounds. In particular, the combinations of the mGluR5 inhibiting compound of this invention can be advantageously used in combination with i) opiate agonists or antagonists, ii) calcium channel antagonists, iii) 5HT receptor agonists or antagonists iv) sodium channel antagonists, v) NMDA receptor agonists or antagonists, vi) COX-2 selective  
25 inhibitors, vii) NK1 antagonists, viii) non-steroidal anti-inflammatory drugs (“NSAID”), ix) GABA-A receptor modulators, x) dopamine agonists or antagonists, xi) selective serotonin reuptake inhibitors (“SSRI”) and/or selective serotonin and norepinephrine reuptake inhibitors (“SSNRI”), xii) tricyclic antidepressant drugs, xiii) norepinephrine modulators, xiv) L-DOPA, xv) buspirone, xvi) lithium, xvii)  
30 valproate, xviii) neurontin (gabapentin), xix) olanzapine, xx) nicotinic agonists or antagonists including nicotine, xxi) muscarinic agonists or antagonists, xxii) heroin substituting drugs such as methadone, levo-alpha-acetylmethadol, buprenorphine and naltrexone, and xxiii) disulfiram and acamprosate.

The abbreviations used herein have the following tabulated meanings. Abbreviations not tabulated below have their meanings as commonly used unless specifically stated otherwise.

Ac	acetyl
AIBN	2,2'-azobis(isobutyronitrile)
BINAP	1,1'-bi-2-naphthol
Bn	benzyl
CAMP	cyclic adenosine-3',5'-monophosphate
DAST	(diethylamino)sulfur trifluoride
DEAD	diethyl azodicarboxylate
DBU	1,8-diazabicyclo[5.4.0]undec-7-ene
DIBAL	diisobutylaluminum hydride
DMAP	4-(dimethylamino)pyridine
DMF	N,N-dimethylformamide
Dppf	1,1'-bis(diphenylphosphino)-ferrocene
EDCI	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
Et <sub>3</sub> N	triethylamine
GST	glutathione transferase
HMDS	hexamethyldisilazide
LDA	lithium diisopropylamide
m-CPBA	metachloroperbenzoic acid
MMPP	monoperoxyphthalic acid
MPPM	monoperoxyphthalic acid, magnesium salt 6H <sub>2</sub> O
Ms	methanesulfonyl = mesyl = SO <sub>2</sub> Me
MsO	methanesulfonate = mesylate
NBS	N-bromo succinimide
NSAID	non-steroidal anti-inflammatory drug
o-Tol	ortho-tolyl
OXONE®	2KHSO <sub>5</sub> •KHSO <sub>4</sub> •K <sub>2</sub> SO <sub>4</sub>
PCC	pyridinium chlorochromate
Pd <sub>2</sub> (dba) <sub>3</sub>	Bis(dibenzylideneacetone) palladium(0)

PDC	pyridinium dichromate
PDE	Phosphodiesterase
Ph	Phenyl
Phe	Benzenediyl
PMB	para-methoxybenzyl
Pye	Pyridinediyl
r.t.	room temperature
Rac.	Racemic
SAM	aminosulfonyl or sulfonamide or SO <sub>2</sub> NH <sub>2</sub>
SEM	2-(trimethylsilyl)ethoxymethoxy
SPA	scintillation proximity assay
TBAF	tetra-n-butylammonium fluoride
Th	2- or 3-thienyl
TFA	trifluoroacetic acid
TFAA	trifluoroacetic acid anhydride
THF	Tetrahydrofuran
Thi	Thiophenediyl
TLC	thin layer chromatography
TMS-CN	trimethylsilyl cyanide
TMSI	trimethylsilyl iodide
Tz	1H (or 2H)-tetrazol-5-yl
XANTPHOS	4,5-Bis-diphenylphosphanyl-9,9-dimethyl-9H-xanthene
C <sub>3</sub> H <sub>5</sub>	Allyl

#### ALKYL GROUP ABBREVIATIONS

Me	=	Methyl
Et	=	ethyl
<i>n</i> -Pr	=	normal propyl
<i>i</i> -Pr	=	isopropyl
<i>n</i> -Bu	=	normal butyl



<i>i</i> -Bu	=	isobutyl
<i>s</i> -Bu	=	secondary butyl
<i>t</i> -Bu	=	tertiary butyl
c-Pr	=	cyclopropyl
c-Bu	=	cyclobutyl
c-Pen	=	cyclopentyl
c-Hex	=	cyclohexyl

### ASSAYS DEMONSTRATING BIOLOGICAL ACTIVITY

The compounds of this invention were tested against the hmGluR5a  
 5 receptor stably expressed in mouse fibroblast Ltk<sup>-</sup> cells (the hmGluR5a/L38-20 cell  
 line) and activity was detected by changes in  $[Ca^{++}]_i$ , measured using the fluorescent  
 $Ca^{++}$ -sensitive dye, fura-2. InsP assays were performed in mouse fibroblast Ltk<sup>-</sup> cells  
 (LM5a cell line) stably expressing hmGluR5a. The assays described in International  
 Patent Publication WO 0116121 can be used.

10

#### Calcium Flux Assay

The activity of compounds was examined against the hmGluR5a  
 receptor stably expressed in mouse fibroblast Ltk<sup>-</sup> cells (the hmGluR5a/L38 cell line).  
 See generally Daggett et al., *Neuropharmacology* 34:871-886 (1995). Receptor  
 15 activity was detected by changes in intracellular calcium ( $[Ca^{2+}]_i$ ) measured using the  
 fluorescent calcium-sensitive dye, fura-2. The hmGluR5a/L38-20 cells were plated  
 onto 96-well plates, and loaded with 3  $\mu$ M fura-2 for 1h. Unincorporated dye was  
 washed from the cells, and the cell plate was transferred to a 96-channel fluorimeter  
 (SIBIA-SAIC, La Jolla, CA) which is integrated into a fully automated plate handling  
 20 and liquid delivery system. Cells were excited at 350 and 385nm with a xenon source  
 combined with optical filters. Emitted light was collected from the sample through a  
 dichroic mirror and a 510nm interference filter and directed into a cooled CCD  
 camera (Princeton Instruments). Image pairs were captured approximately every 1s,  
 and ratio images were generated after background subtraction. After a basal reading  
 25 of 20s, an EC<sub>80</sub> concentration of glutamate (10 $\mu$ M) was added to the well, and the  
 response evaluated for another 60s. The glutamate-evoked increase in  $[Ca]_i$  in the

presence of the screening compound was compared to the response of glutamate alone (the positive control).

#### **Phosphatidylinositol hydrolysis (PI) assays**

- 5                   Inositolphosphate assays were performed as described by Berridge et al. [Berridge et al, *Biochem. J.* 206: 587-5950 (1982); and Nakajima et al., *J. Biol. Chem.* 267:2437-2442 (1992)] with slight modifications. Mouse fibroblast Ltk cells expressing hmGluR5 (hmGluR5/L38- 20 cells) were seeded in 24-well plates at a density of 8x10<sup>5</sup>cells/well. One  $\mu$ Ci of [<sup>3</sup>H]-inositol (Amersham PT6-271; Arlington Heights, Ill.; specific activity = 17.7 Ci/mmol) was added to each well and incubated for 16h at 37°C. Cells were washed twice and incubated for 45min in 0.5mL of standard Hepes buffered saline buffer (HBS; 125mM NaCl, 5mM KCl, 0.62mM MgSO<sub>4</sub>, 1.8mM CaCl<sub>2</sub>, 20mM HEPES, 6mM glucose, pH to 7.4). The cells were washed with HBS containing 10mM LiCl, and 400 $\mu$ L buffer added to each well.
- 10                   Cells were incubated at 37°C for 20min. For testing, 50 $\mu$ L of 10X compounds used in the practice of the invention (made in HBS/LiCl (100mM)) was added and incubated for 10 minutes. Cells were activated by the addition of 10 $\mu$ M glutamate, and the plates left for 1 hour at 37°C. The incubations were terminated by the addition of 1mL ice-cold methanol to each well. In order to isolate inositol
- 15                   phosphates (IPs), the cells were scraped from wells, and placed in numbered glass test tubes. One mL of chloroform was added to each tube, the tubes were mixed, and the phases separated by centrifugation. IPs were separated on Dowex anion exchange columns (AG 1-X8 100-200 mesh formate form). The upper aqueous layer (750 $\mu$ L) was added to the Dowex columns, and the columns eluted with 3mL of distilled
- 20                   water. The eluents were discarded, and the columns were washed with 10mLs of 60mM ammonium formate/5mM Borax, which was also discarded as waste. Finally, the columns were eluted with 4mL of 800mM ammonium formate/0.1M formic acid, and the samples collected in scintillation vials. Scintillant was added to each vial, and the vials shaken, and counted in a scintillation counter after 2 hours.
- 25                   Phosphatidylinositol hydrolysis in cells treated with certain exemplary compounds was compared to phosphatidylinositol hydrolysis in cells treated with the agonist alone in the absence of compound.
- 30                   The compounds of this application have mGluR5 inhibitory activity as shown by IC<sub>50</sub> values of less than 10  $\mu$ M in the calcium flux assay or inhibition at a concentration of 100  $\mu$ M in the PI assay. Preferably, the compounds should have IC<sub>50</sub>

35                   The compounds of this application have mGluR5 inhibitory activity as shown by IC<sub>50</sub> values of less than 10  $\mu$ M in the calcium flux assay or inhibition at a concentration of 100  $\mu$ M in the PI assay. Preferably, the compounds should have IC<sub>50</sub>

values of less than 1  $\mu\text{M}$  in the calcium flux assay and  $\text{IC}_{50}$  values of less than 10  $\mu\text{M}$  in the PI assay. Even more preferably, the compounds should have  $\text{IC}_{50}$  values of less than 500 nM in the calcium flux assay and  $\text{IC}_{50}$  values of less than 1  $\mu\text{M}$  in the PI assay

5                   Examples 1 to 347 have mGluR5 inhibitory activity as shown by inhibition at 10  $\mu\text{M}$  or less in the calcium flux assay or inhibition at 100  $\mu\text{M}$  or less in the PI assay.

                  The examples that follow are intended as an illustration of certain preferred embodiments of the invention and no limitation of the invention is implied.

10                  Unless specifically stated otherwise, the experimental procedures were performed under the following conditions. All operations were carried out at room or ambient temperature - that is, at a temperature in the range of 18-25°C. Evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 pascals: 4.5-30 mm. Hg) with a bath temperature of up to 60°C. The course of

15       reactions was followed by thin layer chromatography (TLC) and reaction times are given for illustration only. Melting points are uncorrected and 'd' indicates decomposition. The melting points given are those obtained for the materials prepared as described. Polymorphism may result in isolation of materials with different melting points in some preparations. The structure and purity of all final

20       products were assured by at least one of the following techniques: TLC, mass spectrometry, nuclear magnetic resonance (NMR) spectrometry or microanalytical data. When given, yields are for illustration only. When given, NMR data is in the form of delta ( $\delta$ ) values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as internal standard, determined at 300 MHz,

25       400 MHz or 500 MHz using the indicated solvent. Conventional abbreviations used for signal shape are: s. singlet; d. doublet; t. triplet; m. multiplet; br. broad; etc. In addition, "Ar" signifies an aromatic signal. Chemical symbols have their usual meanings; the following abbreviations are used: v (volume), w (weight), b.p. (boiling point), m.p. (melting point), L (liter(s)), mL (milliliters), g (gram(s)), mg

30       (milligrams(s)), mol (moles), mmol (millimoles), eq (equivalent(s)).


### Methods of Synthesis

Compounds of the present invention can be prepared according to the following methods. The substituents are the same as in Formula I except where defined otherwise.

In accordance with another embodiment of the present invention, there are provided methods for the preparation of heteroaryl-substituted tetrazole compounds as described above. For example, many of the heterocyclic compounds described above can be prepared using synthetic chemistry techniques well known in the art (see *Comprehensive Heterocyclic Chemistry*, Katritzky, A. R. and Rees, C. W. eds., Pergamon Press, Oxford, 1984) from a heteroaryl-substituted tetrazole of Formula (I).


In Schemes 1 to 3 below, W, X, Y, Z, A and B are as defined above for Formula (I).

#### Scheme 1



Referring to **Scheme 1**, ring system **Y** containing an aldehyde moiety (prepared using synthetic chemistry techniques well known in the art) is reacted with an arylsulfonylhydrazide in a suitable solvent (*e.g.* EtOH, MeOH, THF, DME, DMF *etc.*) at a temperature between 0°C to 100°C for 5 to 60min to form an arylsulfonylhydrazone. An amine-substituted **X** (prepared using synthetic chemistry techniques well known in the art) is treated with nitrous acid, at a temperature of –10°C to 0°C, in a suitable solvent such as, for example, water. In this manner an arenediazonium species is generated which then reacts with an arylsulfonylhydrazone in a 1,3-dipolar cycloaddition reaction to form a substituted tetrazole as shown (for example, see A.S. Shawali et al., *J. Heterocyclic Chem.* **1979**, 16, 123-128). The product from **Scheme 1**, a disubstituted tetrazole, can be isolated and purified employing standard techniques, such as solvent extraction, acid-base extraction, chromatography, crystallization, distillation and the like.


#### 30 Scheme 2



As shown in **Scheme 2** above, **Y** substituted with a nitrile functional group (prepared using methods well known in the art) is reacted with an azide moiety, such as  $\text{LiN}_3$ ,  $\text{NaN}_3$  or  $\text{TMSN}_3$ , in a suitable solvent (*e.g.* toluene, benzene, xylenes *etc.*) at a temperature in the range of about  $25^\circ\text{C}$  to  $180^\circ\text{C}$  to form a monosubstituted tetrazole. This reaction can conveniently be performed with an added catalyst such as dibutyltin oxide. The resulting tetrazole may then be coupled with **X** substituted with a group **T**. **T** maybe a metalloid species such as  $\text{B(OR)}_2$ ,  $\text{BiLn}$  and the like and the reaction may be promoted with stoichiometric or catalytic amounts of metal salts such as  $\text{Cu(OAc)}_2$ ,  $\text{CuI}$  or  $\text{CuOTf}$  and the like. Conveniently, a base (*e.g.* pyridine,  $\text{NEt}_3$ ,  $\text{Cs}_2\text{CO}_3$ ,  $\text{K}_2\text{CO}_3$  *etc.*) will also be present and the reaction is carried out in a suitable solvent (*e.g.* DCM, THF, DME toluene, MeCN, DMF,  $\text{H}_2\text{O}$  *etc.*). Additionally, molecular sieves may be used as a cocatalyst (see for example Fedorov, A. Y.; Finet, J-P. *Tetrahedron Lett.* **1999**, 40, 2747-2748).

Alternatively **T** may be a halogen or other functional group capable of undergoing a metal catalyzed *N*-arylation cross-coupling reaction in which case additional promoters such as 1,10-phenanthroline and dibenzylideneacetone may also be added to the reaction mixture. The cross-coupling reaction maybe carried out at ambient temperature or heated to a temperature between about  $30^\circ\text{C}$  to  $150^\circ\text{C}$ . The reaction mixture is then maintained at a suitable temperature for a time in the range of about 4 to 72 hours, with 18 hours typically being sufficient (see for example Lam, P. Y. S.; Clark, C. G.; Saubern, S.; Adams, J.; Winters, M. P.; Cham, D. M. T.; Combs, A. *Tetrahedron Lett.* **1998**, 39, 2941-2944 and Kiyomori, A.; Marcoux, J. F.; Buchwald, S. L. *Tetrahedron Lett.* **1999**, 40, 2657-2660). The product from **Scheme 2**, a disubstituted tetrazole, can be isolated and purified employing standard techniques, such as solvent extraction, acid-base extraction, chromatography, crystallization, distillation and the like.

### Scheme 3




Referring to **Scheme 3**, the monosubstituted tetrazole is prepared as described in **Scheme 2** from a suitable nitrile-substituted precursor. The tetrazole is then reacted with an *N*-fluoropyridinium salt, which may be optionally substituted, in the presence of a suitable base (*e.g.* MeONa, EtONa, *t*BuOK and the like) for a period of time sufficient for the reaction to proceed to completion, typically from about 1 to 12h, at a temperature in the range of about  $-100^{\circ}\text{C}$  to  $50^{\circ}\text{C}$ , with  $-78^{\circ}\text{C}$  to  $23^{\circ}\text{C}$  being advantageous (see for example Kiselyov, A. S. and Strekowski, L. *J. Heterocyclic Chem.* **1993**, 30, 1361-1364). The product from **Scheme 3**, a 2-pyridyltetrazole derivative, can be isolated and purified employing standard techniques, such as solvent extraction, acid-base extraction, chromatography, crystallization, distillation and the like.

In the schemes above, ring systems X and/or Y may already contain a pendant ring W and/or Z. However, if required, ring systems W and/or Z may be appended to X and/or Y respectively where G and/or J are functional groups capable of undergoing a metal catalyzed-cross coupling (such as halogen, trifluoromethanesulfonate,  $\text{B(OR)}_2$ ,  $\text{ZnX}$ ,  $\text{SnR}_3$ , and the like -**Scheme 4** below). Ring systems W and Z are substituted with groups P, Q, S and T which may be for example, halogen, trifluoromethanesulfonate,  $\text{B(OR)}_2$ ,  $\text{ZnX}$ ,  $\text{SnR}_3$ , and the like. Typically, a transition metal catalyst such as  $\text{Pd(PPh}_3)_4$ ,  $\text{Pd(PPh}_3)_2\text{Cl}_2$ ,  $\text{Pd(OAc)}_2$ ,  $\text{NiCl}_2(\text{dppe})$ ,  $\text{Pd(OAc)}_2$ ,  $\text{Pd}_2(\text{dba})_3$ ,  $\text{Cu(OAc)}_2$ , CuI or the like may be employed, typically along with a suitable base such as  $\text{K}_2\text{CO}_3$ ,  $\text{K}_3\text{PO}_4$ ,  $\text{Cs}_2\text{CO}_3$ ,  $\text{Et}_3\text{N}$ , pyridine or the like. Additionally, ligands such as BINAP, di-*tert*-butyl phosphinobiphenyl, di-cyclohexylphosphino biphenyl, tri *tert*-butylphosphine, XANTPHOS, triphenylarsine and the like may be added. The reaction is carried out in a suitable solvent such as toluene, DME, dioxane, THF, water or a combination of the above and is typically heated at  $50^{\circ}\text{C}$  –  $150^{\circ}\text{C}$  for between 1 and 48 hrs. The reaction may be homogeneous or heterogeneous (see for example Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, 95, 2457-2483 and Dai, C.; Fu, G.C *J. Am. Chem. Soc.*, **2001**, 123, 2719-2724 and Littke, A.F.; Fu, G.C. *Angew.*

*Chem. Int. Ed.* **1999**, 38, 6, 2411-2413 and Dai, C; Fu, G.C. *J. Am. Chem. Soc.* **2001**, 123, 2719-2724).

**Scheme 4**



5

Alternatively ring systems W or Z may be a nitrogen containing heterocycle wherein the nitrogen is directly attached to the ring system X or Y respectively. In this case G and/or J are groups capable of undergoing a metal catalyzed N-aryl cross-coupling (such as halogen, trifluoromethane-sulfonate, B(OR)<sub>2</sub>, ZnX, SnR<sub>3</sub>, and the like –

- 10 **Scheme 4**). Typically a transition metal such as CuI, Cu(OAc)<sub>2</sub>, Cu(OTf)<sub>2</sub>, Pd(PPh<sub>3</sub>)<sub>4</sub>, Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub>, Pd(OAc)<sub>2</sub>, Pd<sub>2</sub>(dba)<sub>3</sub>, NiCl<sub>2</sub>(dppe) is used along with a suitable base such as K<sub>2</sub>CO<sub>3</sub>, K<sub>3</sub>PO<sub>4</sub>, Cs<sub>2</sub>CO<sub>3</sub>, NaOtBu or the like. Additionally, phosphine containing ligands such as BINAP, di-*tert*-butyl phosphinobiphenyl, di-cyclohexylphosphino biphenyl, tri *tert*-butylphosphine, XANTPHOS and the like may
- 15 be added. Further, additives such as 1,10-phenanthroline, 1,2-diaminocyclohexane, dibenzylideneacetone may be used. The reaction is typically carried out in a solvent such as toluene, DME, dioxane, THF, water or a combination of the above and is typically heated at 50°C – 150°C for between 1 and 48 hrs. The reaction may be homogeneous or heterogeneous. The product from **Scheme 4**, can be isolated and
- 20 purified employing standard techniques, such as solvent extraction, acid-base extraction, chromatography, crystallization, distillation and the like (see for example Lam, P. Y. S.; Clark, C. G.; Saubern, S.; Adams, J.; Winters, M. P.; Cham, D. M. T.; Combs, A. *Tetrahedron Lett.* **1998**, 39, 2941-2944 and Kiyomori, A.; Marcoux, J. F.;

Buchwald, S. L. *Tetrahedron Lett.* **1999**, *40*, 2657-2660 and Wolfe, J.P.; Tomori, H.; Sadighi, J.P.; Yin, J.; Buchwald, S.L. *J. Org. Chem.*, **2000**, *65*, 1158-1174 and Yin, J.; Buchwald, S.L.; *Org. Lett.*, **2000**, *2*, 1101-1104).

In addition, many of the heterocyclic intermediate compounds described above can be prepared using other synthetic chemistry techniques well known in the art (see *Comprehensive Heterocyclic Chemistry*, Katritzky, A. R. and Rees, C. W. eds., Pergamon Press, Oxford, 1984) and references cited there within.

### EXAMPLE 1

#### 10      **1-Methyl-3-[3-(5-pyridin-2-yl-2H-tetrazol-2-yl)phenyl]imidazolidin-2-one**

3-(3-Methylimidazolidin-2-one)aniline (76mg, 0.4mmol) was dissolved in 6N HCl (0.5mL, 3.0mmol), cooled to 0°C, and a solution of NaNO<sub>2</sub> (30mg, 0.43mmol) in H<sub>2</sub>O (0.5mL) was added dropwise. The internal reaction temperature was maintained at <5°C by the addition of ice chips to the flask.

15 Separately, 2-pyridyl carboxaldehyde (43mg, 0.4mmol) and toluenesulfonyl hydrazide (69mg, 0.4mmol) were combined in ethanol (0.5mL). The resulting reaction mixture was stirred at ambient temperature for 15min and monitored by TLC for the disappearance of aldehyde (2,4-DNP stain). NaOH pellets (160mg, 4.0mmol) were then added along with H<sub>2</sub>O (1mL) and the reaction mixture was cooled to 0°C with an  
20 ice bath.

The 3-(3-methylimidazolidin-2-one)aniline diazotization reaction contents were then added dropwise *via* pipet. The resulting reaction was stirred for an additional 10 min at 0°C, then the cooling bath was removed and the reaction was allowed to warm to ambient temperature. The crude mixture was diluted with EtOAc  
25 (10mL) and washed with H<sub>2</sub>O (3x10mL). The organic phase was dried (MgSO<sub>4</sub>), filtered, and concentrated *in vacuo*. The residue was chromatographed on silica gel eluting with hexanes:EtOAc (3:1) to afford 2-[2-(3-(3-methylimidazolidin-2-one)phenyl)-2H-tetrazol-5-yl]pyridine as a pale orange solid. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz) δ 8.84 (d, 1H), 8.33 (d, 1H), 8.20-8.21 (m, 1H), 7.98-8.02 (m, 1H), 7.87-7.92  
30 (m, 2H), 7.41-7.54 (m, 2H), 3.89-3.95 (m, 2H), 3.52-3.57 (m, 2H), 2.93 (s, 3H). MS (ESI) 322.1 (M+H)<sup>+</sup>.

### COMPOUND 1

#### **2-[2-(4-Bromophenyl)-2H-tetrazol-5-yl]pyridine**



Following the procedure described in **EXAMPLE 1** for the synthesis of 1-methyl-3-[3-(5-pyridin-2-yl-2*H*-tetrazol-2-yl)phenyl]imidazolidin-2-one, but using 4-bromoaniline (17.2 g, 100mmol) and pyridine-2-carboxaldehyde (10.7 g, 100mmol), 2-[2-(4-bromophenyl)-2*H*-tetrazol-5-yl]pyridine was obtained as an orange solid. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz) δ 8.83 (d, 1H), 8.33 (d, 1H), 8.15 (d, 2H), 7.96-7.87 (m, 1H), 7.70 (d, 2H), 7.44 (dd, 1H).

### **EXAMPLE 2**

#### **2-[2-(4-Pyridin-2-ylphenyl)-2*H*-tetrazol-5-yl]pyridine**

10 A solution of 2-[2-(4-bromophenyl)-2*H*-tetrazol-5-yl]pyridine (300mg, 1mmol), 2-pyridylzinc bromide (0.5M in THF, 2 mL) and tetrakis-(triphenylphosphine)palladium(0) (100 mg, 0.085 mmol) in anhydrous THF (10 mL) was degassed for 15 min and heated to 60°C. After 4 hours, a further equivalent of 2-pyridylzinc bromide (0.5M in THF, 2 mL) was added and the reaction mixture was  
15 heated at reflux overnight. The reaction mixture was then quenched with brine (25 mL), extracted with EtOAc (3x25 mL), dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. The crude residue was chromatographed on silica gel, eluting with EtOAc to afford a yellow solid. This was dissolved in anhydrous THF (10 mL) and HCl in ether (1.5 equiv) was added and the resulting precipitate was filtered to afford a yellow solid.  
20 <sup>1</sup>H-NMR (CD<sub>3</sub>OD, 300 MHz) δ 9.00-8.94 (m, 2H), 8.81-8.70 (m, 2H), 8.67-8.61 (m, 3H), 8.53 (d, 1H), 8.33 (d, 2H), 8.17-8.09 (m, 2H). MS (ESI) 301.1 (M+H)<sup>+</sup>.

### **EXAMPLE 3**

#### **2-[2-(4-Pyridin-4-ylphenyl)-2*H*-tetrazol-5-yl]pyridine**

25 A solution of 2-[2-(4-bromophenyl)-2*H*-tetrazol-5-yl]pyridine (300mg, 1mmol), potassium carbonate (0.276g, 2.0mmol), and pyridin-4-ylboronic acid (123mg, 1mmol) in ethylene glycol dimethyl ether (20mL) and water (4mL) was degassed with argon for 15min. Then tetrakis(triphenylphosphine)palladium(0) (80mg, 0.068mmol) was added, the mixture degassed for another 15min, and the  
30 solution was stirred at 65°C for 12h. The reaction mixture was then quenched with water (30mL), extracted with EtOAc (3 X 30mL) and washed with brine. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. The crude residue was chromatographed on silica gel, eluting with EtOAc to afford a yellow solid. This was dissolved in anhydrous THF (10mL) and HCl (1.5equiv) in ether was added. The

resulting precipitate was filtered off to afford 2-[2-(4-pyridin-4-ylphenyl)-2*H*-tetrazol-5-yl]pyridine as a yellow solid. <sup>1</sup>H NMR (CD<sub>3</sub>OD, 300 MHz) δ 8.99-8.97 (m, 3H), 8.74 (d, 1H), 8.62-8.54 (m, 5H), 8.35 (d, 2H), 8.05 (t, 1H). MS (ESI) 301.0 (M+H)<sup>+</sup>.

5

### **COMPOUND 2**

#### **2-[2-(3-Iodophenyl)-2*H*-tetrazol-5-yl]pyridine**

Following the procedure described in EXAMPLE 1 for the synthesis of 1-methyl-3-[3-(5-pyridin-2-yl-2*H*-tetrazol-2-yl)phenyl]imidazolidin-2-one, but using 3-iodoaniline (11g, 50mmol) and pyridine-2-carboxaldehyde (4.7mL, 50mmol), 2-[2-(3-iodophenyl)-2*H*-tetrazol-5-yl]pyridine was obtained as a biege solid. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz) δ NMR 8.92 (d, 1H), 8.69 (m, 1H), 8.44 (d, 1H), 8.30 (d, 1H), 8.03 – 8.06 (m, 1H), 7.87 – 7.88 (m, 1H), 7.56 – 7.59 (m, 1H), 7.32 – 7.35 (m, 1H).

15

### **EXAMPLE 4**

#### **2-[2-[3-(1*H*-Imidazol-1-yl)phenyl]-2*H*-tetrazol-5-yl]pyridine**

2-[2-(3-Iodophenyl)-2*H*-tetrazol-5-yl]pyridine (520mg, 1.5mmol), imidazole (133mg, 2.0mmol), potassium carbonate (435mg, 3.2mmol), copper (I) iodide (57mg, 0.3mmol) and 1,2-diamino cyclohexane (90μL, 0.8mmol) were weighed into a flask and flushed with Ar(g). Dry dioxane (0.7mL) was added and the reaction mixture heated at 100°C for 18h. The reaction mixture was then poured into NH<sub>4</sub>OH (10% in H<sub>2</sub>O), NH<sub>4</sub>Cl (sat aq) (1/1, 10mL) and shaken with EtOAc (30mL). The organic layer was separated and the aqueous layer extracted with EtOAc (3 x 30mL), the combined organics were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated onto silica gel. Purification by liquid chromatography on silica gel (hexane:EtOAc = 90:10) gave a yellow solid. This was recrystallized from EtOAc/Hexane to give yellow crystals which were dissolved in CH<sub>2</sub>Cl<sub>2</sub>/Et<sub>2</sub>O and HCl in ether (1.5equiv) was added. The resulting precipitate was filtered off to afford 2-{2-[3-(1*H*-imidazol-1-yl)phenyl]-2*H*-tetrazol-5-yl}pyridine as a yellow solid. <sup>1</sup>H NMR ((CD<sub>3</sub>)<sub>2</sub>SO, 300 MHz) δ 9.94 (s, 1H), 8.23 – 8.84 (m, 1H), 8.68 – 8.69 (m, 1H), 8.48 – 8.49 (m, 1H), 8.39 – 8.42 (m, 1H), 8.30 – 8.33 (m, 1H) 8.07 – 8.13 (m, 2H), 7.97 – 8.03 (m, 2H), 7.63 – 7.67 (m, 1H). MS (ESI) 262.2 (MH-N<sub>2</sub>)<sup>+</sup>.

25

30

### **COMPOUND 3**

#### **2-(2-Methoxy-4-nitrophenyl)pyridine**

2-Pyridylzinc bromide (15.0mmol, 30.0mL of 0.5M solution in THF) was added via syringe to a stirred solution of 2-iodo-5-nitroanisole (4.0g, 14.3mmol) and tetrakis(triphenylphosphine)palladium (828mg, 0.7mmol) in deoxygenated THF. The mixture was stirred at 60°C for 16h, followed by the addition of additional  
5 tetrakis(triphenylphosphine)palladium (170mg, 0.14mmol) and 2-pyridylzinc bromide (5.0mmol, 10.0mL of 0.5M solution in THF). Continued stirring at 60°C resulted in the disappearance by TLC analysis of the starting material 2-iodo-5-nitroanisole. The reaction mixture was concentrated *in vacuo*, dissolved in EtOAc (200mL) and partitioned with H<sub>2</sub>O (100mL). The EtOAc was washed with an additional portion of  
10 H<sub>2</sub>O, then dried (MgSO<sub>4</sub>), filtered, and concentrated *in vacuo*. The crude residue was purified by silica gel chromatography, eluting with 3:1 hexanes:EtOAc to afford 2-(2-methoxy-4-nitrophenyl) as a white solid. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz) δ 8.75 (m, 1H), 7.95-7.98 (m, 2H), 7.86-7.91 (m, 2H), 7.75-7.80 (m, 1H), 7.28-7.33 (m, 1H), 3.98 (s, 3H).

15

#### COMPOUND 4

##### **3-Methoxy-4-pyridin-2-ylaniline**

Tin(II) chloride dihydrate (6.77g, 30.0mmol) and 2-(2-methoxy-4-nitrophenyl)pyridine (2.30g, 10.0mmol) were combined in MeOH (100mL) and  
20 stirred at reflux for 18h. The reaction mixture was then cooled to RT and concentrated *in vacuo*. The residue was then partitioned with EtOAc (200mL) and 1N aqueous NaOH (100mL). The aqueous layer was washed with EtOAc (3x75 mL), and the combined EtOAc layers were back-extracted with H<sub>2</sub>O (100mL), dried (MgSO<sub>4</sub>), filtered, and concentrated *in vacuo*. The resulting residue was purified by SiO<sub>2</sub>  
25 chromatography, eluting with 2:1 EtOAc:hexanes to afford 3-methoxy-4-pyridin-2-ylaniline as an amber oil. <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz) δ 8.63 (m, 1H), 7.78 (d, 1H), 7.57-7.67 (m, 2H), 7.04-7.13 (m, 1H), 6.34-6.41(dd, 1H), 6.26 (d, 1H), 3.72-3.92 (s, 2H), 3.72 (s, 3H).

30

#### COMPOUND 5

##### **3-Methoxy-4-(3-methyl-1,2,4-oxadiazol-5-yl)aniline**

To a suspension of methyl amidoxime hydrochloride (1.8g, 16.3mmol) in THF (80mL) at rt was added NaH (1.32g of a 60% dispersion in oil, 33.1mmol). Reaction was stirred at 60°C for 15min, then methyl-4-amino-2-methoxy benzoate  
35 (1.0g, 5.5mmol) was added and stirring was continued. After 4h, DMF (60mL) was

added to aid solubility and the reaction temperature was increased to 90°C. After 1.5h, reaction was complete by TLC analysis. Reaction was quenched at rt with H<sub>2</sub>O (20mL), then concentrated via rotary evaporation. The crude residue was partitioned in a separatory funnel with EtOAc (100mL) and H<sub>2</sub>O (50mL). The EtOAc layer was washed with an additional portion of H<sub>2</sub>O (50mL), then dried (MgSO<sub>4</sub>), filtered, and concentrated *in vacuo*. The residue was then purified by silica gel chromatography, eluting with 3:1 EtOAc:Hexanes to afford 3-methoxy-4-(3-methyl-1,2,4-oxadiazol-5-yl)aniline as a tan solid. <sup>1</sup>H NMR (CD<sub>3</sub>OD, 500 MHz) δ 7.72 (m, 1H), 6.31-6.37 (m, 2H), 3.87 (s, 3H), 2.36 (s, 3H). MS (ESI) 206.1 (M+H)<sup>+</sup>.

10

### COMPOUND 6

#### **2-{2-[3-(Trimethylstannyl)phenyl]-2H-tetrazol-5-yl}pyridine**

A solution of 2-[2-(3-bromophenyl)-2H-tetrazol-5-yl]pyridine (750mg, 2.5mmol), hexamethylditin (895mg, 2.8mmol), cesium fluoride (418mg, 2.8mmol) and tetrakis(triphenylphosphine)palladium(0) (145mg, 0.13mmol) in 20mL of dry toluene was degassed with nitrogen for 15min. The solution was heated to 100°C and stirred under nitrogen for 6h. The reaction mixture was allowed to cool to rt, diluted with 10mL of ethyl acetate, and then filtered through a pad of Celite. Water (40mL) was added to the solution, extracted with 3 X 40mL of ethyl acetate, and then the organic extracts were combined and washed with brine. The extracts were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and then concentrated. The crude material was chromatographed over silica gel, eluting with 30% EtOAc/hexanes. Purification yielded a yellow oil, which crystallized upon sitting to afford 2-{2-[3-(trimethylstannyl)phenyl]-2H-tetrazol-5-yl}pyridine as a light yellow solid. <sup>1</sup>H NMR (CD<sub>3</sub>OD, 500 MHz) δ 8.72-8.88 (m, 1H), 8.39-8.43 (m, 2H), 8.21-8.22 (m, 1H), 7.92-7.95 (m, 1H), 7.60-7.62 (m, 1H), 7.52-7.55 (m, 1H), 7.46-7.48 (m, 1H), 0.40 (s, 9H).

25

### EXAMPLE 5

#### **2-[2-(2-Pyrazin-3-ylphenyl)-2H-tetrazol-5-yl]pyridine**

A vial containing 2-[2-(3-trimethylstannylylphenyl)-2H-tetrazol-5-yl]pyridine (70mg, 0.18mmol), chloropyrazine (23mg, 2.0mmol), tris(dibenzylideneacetone)dipalladium(0) (10mg, 0.01mmol), cesium fluoride (60mg, 0.4mmol), and tri-*t*-butyl-phosphine (0.101mL of 10% weight in hexanes, 0.05mmol) was sealed with a septum cap and purged with nitrogen. Degassed dioxane (2mL) was added to the vial via syringe and stirred under nitrogen for 10min. The reaction

35

mixture was subjected to microwave conditions at 140°C for 600s using the Personal Chemistry Smith Creator microwave instrument. The reaction mixture was diluted with EtOAc, then extracted with 3X EtOAc and water. The organic extracts were combined, dried over NaSO<sub>4</sub>, filtered, and then concentrated. The crude material was chromatographed over silica gel, eluting with 75% EtOAc/hexanes, to yield a yellow solid. The solid was further purified with a Waters preparatory HPLC instrument to yield 2-[2-(2-pyrazin-3-ylphenyl)-2*H*-tetrazol-5-yl]pyridine as a light yellow solid. <sup>1</sup>H NMR (CD<sub>3</sub>OD, 500 MHz) 9.34 (s, 1H), 9.08 (s, 1H), 8.98 (m, 1H), 8.86 (s, 1H), 8.81 (m, 1H), 8.69 (m, 1H), 8.62 (t, 1H), 8.41-8.45 (m, 2H), 8.08 (m, 1H), 7.90 (t, 1H). MS 302 (M+H)<sup>+</sup>.

#### EXAMPLE 6

##### 2-[2-(4-Morpholin-3-ylphenyl)-2*H*-tetrazol-5-yl]pyridine

To a flask containing 2-[2-(4-bromophenyl)-2*H*-tetrazol-5-yl]pyridine (150mg, 0.5mmol), tris(dibenzylideneacetone)dipalladium(0) (26mg, 0.025mmol), 2-(dicyclohexylphosphino)biphenyl (18mg, 0.05mmol), and sodium *t*-butoxide (67mg, 0.7mmol), degassed toluene was added and the resulting mixture was stirred under nitrogen until homogeneity. Morpholine (53μL, 0.6mmol) was added to the mixture *via* syringe, and the reaction was heated to 80°C. The solution was then stirred for 18h under nitrogen. The reaction mixture was quenched with water (20mL), extracted with EtOAc (3 X 20mL), and washed with brine. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated *in vacuo*. The crude residue was chromatographed on silica gel eluting with 40% EtOAc in hexanes to afford a yellow solid. This was dissolved in anhydrous THF (10mL) and HCl (1.5equiv) in ether was added, the precipitate was filtered to afford 2-[2-(4-morpholin-3-ylphenyl)-2*H*-tetrazol-5-yl]pyridine as a yellow solid. <sup>1</sup>H NMR ((CD<sub>3</sub>)<sub>2</sub>SO, 300 MHz) δ 8.82 (d, 1H), 8.28 (d, 1H), 8.08 (dt, 1H), 7.51-7.68 (m, 4H), 7.23 (m, 1H), 3.77 (m, 4H), 3.25 (m, 4H). MS (ESI) MS 309 (M+H)<sup>+</sup>.

#### EXAMPLE 7

##### 2-{2-[3-(2*H*-Tetrazol-5-yl)phenyl]-2*H*-tetrazol-5-yl}pyridine

3-(5-Pyridin-2-yl-2*H*-tetrazol-2-yl)benzonitrile (300mg, 1.21mmol), sodium azide (86.5mg, 1.33mmol) and zinc bromide (272mg, 1.21mmol) were suspended in water (3mL). The reaction was capped with a reflux condenser and


stirred at 90°C for 24h, after which time it was cooled to rt and HCl 3N (1.5mL) and ethyl acetate (20mL) were added and the mixture stirred vigorously until no solid was left. The organic layer was isolated and aqueous layer extracted with ethyl acetate (2 x 20mL). The combined organic layers were evaporated, 10mL of 0.25N NaOH was added, the mixture was stirred for 30min, until the original precipitate was dissolved and a suspension of zinc hydroxide was formed. The suspension was filtered, and the solid washed with 10 ml 1 N NaOH. To the filtrate was added 2mL of 3N HCl with stirring causing the tetrazole to precipitate. The tetrazole was filtered and washed with 5mL of 3N HCl and dried in vacuo to afford white powder. <sup>1</sup>H NMR (CD<sub>3</sub>OD, 500 MHz) δ 8.91 (s, 1H), 8.83 (d, 1H), 8.41 (d, 1H), 8.32-8.25 (m, 2H), 8.09 (t, 1H), 7.95 (t, 1H), 7.64 (t, 1H). MS 292.1 (M+H)<sup>+</sup>.









### EXAMPLE 8







#### 2-Pyridin-2-yl-5-(5-pyridin-2-yl-2H-tetrazol-2-yl)benzonitrile

2-[2-(3-Fluoro-4-pyridin-2-ylphenyl)-2H-tetrazol-5-yl]pyridine (300 mg, 0.94 mmol) and sodium cyanide (92 mg, 1.9 mmol) were suspended in methyl sulfoxide (6 ml). The reaction was capped with a reflux condenser and stirred at 170°C for 16 h, after which time it was cooled to ambient temperature. To the reaction solution added ethyl acetate (10 ml) then water (60 ml) causing 2-pyridin-2-yl-5-(5-pyridin-2-yl-2H-tetrazol-2-yl)benzonitrile to precipitate as white powder. <sup>1</sup>H NMR (CD<sub>3</sub>OD, 500 MHz) δ 8.84-8.81 (m, 2H), 8.73 (d, 1H), 8.60 (dd 1H), 8.32 (d, 1H), 8.24 (d, 1H), 8.09-8.07 (m, 2H), 8.01 (d, 1H), 7.64 (m, 1H), 7.58(m, 1H). MS 326.2 (M+H)<sup>+</sup>.








EXAMPLE 9 to EXAMPLE 347 shown below were all prepared similarly to the schemes and procedures described above (ND = not determined).









EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
9.		9.72 (m, 2H), 9.45 (d, 1H), 8.90-8.99 (m, 3H), 8.77 (d, 1H), 8.55-8.62 (m, 2H), 8.32 (dd, 1H), 8.07 (dd, 1H).	MS 302.1 (M+H) <sup>+</sup> .









EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
10.		8.87 (d, 1H), 8.38 (d, 1H), 7.98-7.92 (m, 2H), 7.88 (d, 1H), 7.59-7.38 (m, 7H), 3.97 (s, 3H).	MS 330.6 (M+H) <sup>+</sup> .
11.		8.75-8.77 (d, 1H), 8.66-8.68 (d, 1H), 8.35-8.38 (d, 1H), 8.23 (s, 1H), 8.15-8.19 (d, 1H), 8.06-8.10 (t, 1H), 7.96-8.00 (t, 1H), 7.58-7.62 (m, 3H), 7.46-7.50 (m, 1H), 2.45 (s, 3H)	MS 315.1 (M+H) <sup>+</sup> .
12.		ND	MS 300.1 (M+H) <sup>+</sup> .
13.		ND	MS 300.2 (M+H) <sup>+</sup> .
14.		8.82 – 8.83 (m, 1H), 8.76 (d, 1H), 8.67 (t, 1H), 8.30 (d, 1H), 8.06 – 8.14 (m, 3H), 7.81 – 7.86 (m, 2H), 7.64 (ddd, 1H), 6.64 (dd, 1H).	MS 290.2 (M+H) <sup>+</sup> .
15.		8.83 (d, 1H), 8.50 (s, 1H), 8.32 (d, 1H), 8.18-8.20 (m, 1H), 8.08-8.11 (m, 1H), 7.97 (d, 1H), 7.81 (t, 1H), 7.64 (m, 1H), 7.47 (t, 1H), 7.39 (d, 1H), 7.34 (m, 1H), 7.05 (dd, 1H), 3.92 (s, 3H).	MS 330 (M+H) <sup>+</sup> .
16.		9.32 (s, 2H), 9.29 (s, 1H), 8.83 (m, 1H), 8.59 (t, 1H), 8.28-8.33 (m, 2H), 8.07-8.13 (m, 2H), 7.90 (t, 1H), 7.62 (m, 1H).	MS 302 (M+H) <sup>+</sup> .
17.		8.83 (m, 1H), 8.41 (m, 1H), 8.31 (d, 1H), 8.25-8.27 (m, 1H), 8.10 (dt, 1H), 7.95 (d, 1H), 7.86 (t, 1H), 7.82	MS 315 (M+H) <sup>+</sup> .







EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
		(d, 1H), 7.76 (m, 1H), 7.63-7.66 (m, 2H), 7.40 (d, 1H), 3.90 (brm).	
18.		8.92 – 8.93 (m, 1H), 8.80 – 8.82 (m, 1H), 8.75 – 8.77 (m, 1H), 8.23 – 8.34 (m, 3H), 8.14 – 8.17 (m, 1H), 8.07 (ddd, 1H), 7.97 (ddd, 1H), 7.82 (dd, 1H), 7.62 (ddd, 1H), 7.44 – 7.48 (m, 1H).	MS 301 (M+H) <sup>+</sup> .
19.		9.04 (d, 1H, J = 2.8 Hz), 8.81 – 8.30 (m, 1H), 8.67 (dd, 1H), 8.47 – 8.48 (m, 1H), 8.30 – 8.32 (m, 1H), 8.23 – 8.28 (m, 2H), 8.08 (ddd), 8.02 – 8.04 (m, 1H), 7.85 (dd, 1H), 7.63 (ddd, 1H), 7.55 – 7.60 (m, 1H).	MS 301 (M+H) <sup>+</sup> .
20.		8.81 – 8.83 (m, 1H), 8.71 – 8.73 (m, 2H), 8.51 – 8.52 (m, 1H), 8.26 – 8.32 (m, 2H), 8.05 – 8.11 (m, 2H), 7.84 – 7.89 (m, 3H), 7.62 (ddd, 1H).	MS 301 (M+H) <sup>+</sup> .
21.		8.83 (m, 1H), 8.47 (m, 1H), 8.31 (m, 1H), 8.18 (t, 1H), 8.09 (m, 1H), 8.08(t, 1H), 8.01 (m, 1H), 7.72-7.79 (m, 3H), 7.63 (m, 1H).	MS 306 (M+H) <sup>+</sup>
22.		8.84 (d, 1H), 8.39 (m, 1H), 8.32 (d, 1H), 8.12 (d, 1H), 8.08 (dt, 1H), 7.96 (d, 1H), 7.77 (d, 1H), 7.76 (d, 1H), 7.71 (d, 1H), 7.64 (dd, 1H), 7.24 (dd, 1H)	MS 306 (M+H) <sup>+</sup>
23.		9.10 (s, 2H), 8.83 (m, 1H), 8.51 (m, 1H), 8.31 (d, 1H), 8.20 (m, 1H), 8.12 (dt, 1H), 8.06 (m, 1H), 7.86 (t, 1H), 7.58-7.68 (m, 2H), 4.05 (s, 3H).	MS 332 (M+H) <sup>+</sup>


















EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
24.		8.82 (m, 1H), 8.65 (d, 1H), 8.41 (t, 1H), 8.31 (d, 1H), 8.16-8.21 (m, 2H), 8.09 (dt, 1H), 7.96 (d, 1H), 7.81 (t, 1H), 7.64 (m, 1H), 6.99 (d, 1H), 3.93 (s, 3H).	MS 331 (M+H) <sup>+</sup>
25.		8.47 (d, 1H), 8.38 (d, 1H), 7.98 (s, 1H), 7.91-7.82 (m, 3H), 7.70-7.65 (t, 1H), 7.57-7.53 (m, 1H), 7.34-7.31 (t, 1H), 7.22-7.20 (m, 1H), 6.26 (b, 1H)	MS 357.7 (M+Na) <sup>+</sup>
26.		8.92 (d, 1H), 8.84 (d, 1H), 8.74 (dd, 1H), 8.50 (d, 1H), 8.39 (d, 1H), 8.09-8.27 (m, 4H), 8.01 (d, 1H), 7.73-7.78 (m, 1H), 4.14 (s, 3H).	MS 331.0 (M+H) <sup>+</sup>
27.		8.97 (d, 1H), 8.80 (d, 1H), 8.66 (dd, 1H), 8.52 (d, 1H), 8.18-8.21 (m, 2H), 8.09-8.14 (m, 2H), 8.03 (d, 1H), 4.30 (s, 3H).	MS 337.0 (M+H) <sup>+</sup>
28.		9.42 (s, 1H), 9.11 (d, 1H), 8.99 (d, 1H), 8.91 (br, 1H), 8.72 (s, 1H), 8.70~8.60 (m, 1H), 8.50~8.55 (m, 1H), 8.44 (dd, 1H), 8.28 (dd, 1H), 8.21 (m, 1H), 7.88~7.84 (m, 1H).	MS 335.1 (M+H) <sup>+</sup>
29.		8.24 (s, 1H), 8.94 (dd, 1H), 8.90~8.80 (m, 1H), 8.55 (d, 1H), 8.45~8.8.33 (m, 3H), 8.23 (dd, 1H), 8.78~8.82 (m, 1H), 7.55 (d, 1H), 4.03 (s, 3H).	MS 331.0 (M+H) <sup>+</sup>
30.		ND	MS 290.0 (M+H) <sup>+</sup>







EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
31.		8.84-8.86 (d, 1H), 8.34-8.37 (m, 2H), 8.14-8.18 (m, 1H), 7.89-7.95 (m, 1H), 7.43-7.47 (m, 1H), 7.18-7.21 (d, 1H), 3.91-3.94 (t, 4H), 3.15-3.18 (t, 4H).	MS 343.0 (M+H) <sup>+</sup> .
32.		9.37 (s, 1H) 9.07 (d, 1H), 8.96 (br, 2H), 8.78-8.75 (m, 1H), 8.60 (t, 1H), 8.46 (s, 1H), 8.24-8.23 (m, 2H), 8.13-8.04 (m, 2H).	MS 335.0 (M+H) <sup>+</sup>
33.		8.87 (d, 1H), 8.38 (d, 1H), 7.98-7.92 (m, 2H), 7.88 (d, 1H), 7.59-7.38 (m, 7H), 3.97 (s, 3H).	MS 330.6 (M+H) <sup>+</sup> .
34.		9.05 (m, 2H), 8.84 (m, 2H), 8.75 (1H), 8.46 (m, 3H), 8.219 (m, 3H).	MS 318.8 (M+H) <sup>+</sup> .
35.		8.86 (d, 1H), 8.75 (s, 1H), 8.69 (s, 1H), 8.49 (d, 1H), 8.38 (d, 1H), 7.93 (t, 1H), 7.80 (m, 2H), 7.57 (d, 1H), 7.47 (t, 1H), 7.32 (t, 1H).	MS 359.0 (M+H) <sup>+</sup> .
36.		8.82 (d, 1H), 8.4 (s, 1H), 8.33 (d, 1H), 8.20 (m, 1H), 8.1 (m, 1H), 7.97 (d, 1H), 7.81 (t, 1H), 7.62 (dd, 1H), 7.48 (t, 1H), 7.38 (d, 1H), 7.34 (m, 1H), 7.05 (dd, 1H), 3.85 (s, 3H).	MS 330 (M+H) <sup>+</sup>
37.		9.20 (s, 1H), 9.02 (d, 2H), 8.84 (d, 1H), 8.64 (d, 1H), 8.38 (dd, 1H), 8.33 (d, 1H), 8.11 (dt, 1H), 7.89 (t, 1H), 7.65 (dd, 1H), 7.58 (t, 1H).	MS 302 (M+H) <sup>+</sup>
38.		8.83 (d, 1H), 8.52 (m, 1H), 8.38 (m, 1H), 8.32 (d, 1H), 8.25 (m, 1H), 8.20 (m, 1H), 8.10 (dt, 1H), 8.05 (d, 1H), 7.94 (d, 1H), 7.85 (t, 1H), 7.76 (t, 1H), 7.64 (m, 1H).	MS 325 (M+H) <sup>+</sup>







EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
39.		8.84 (d, 1H), 8.74 (d, 1H), 8.32 (m, 2H), 8.21 (d, 1H), 8.10 (m, 1H), 8.06 (m, 1H), 7.96 (m, 1H), 7.87 (m, 1H), 7.64 (m, 1H).	MS 307 (M+H) <sup>+</sup>
40.		8.83 (d, 1H), 8.40 (s, 1H), 8.32 (d, 1H), 8.19 (m, 1H), 8.10 (dt, 1H), 7.97 (d, 1H), 7.81 (t, 1H), 7.64 (dd, 1H), 7.47 (t, 1H), 7.38 (d, 1H), 7.34 (m, 1H), 7.05 (dd, 1H), 3.88, (s, 3H).	MS 330 (M+H) <sup>+</sup>
41.		8.82 (d, 1H), 8.25-8.32 (m, 3H), 8.09 (dt, 1H), 7.85 (t, 1H), 7.79 (d, 1H), 7.64 (m, 1H), 7.47 (m, 2H), 7.41 (m, 1H), 7.34 (m, 1H), 3.73 (bm, 2H).	MS 315 (M+H) <sup>+</sup>
42.		8.84 (d, 1H), 8.30-8.32 (m, 3H), 8.10 (m, 1H), 7.87 (d, 2H), 7.65 (dd, 1H), 7.44-7.49 (m, 3H), 7.36 (m, 1H).	MS 315 (M+H) <sup>+</sup>
43.		9.06 (d, 1H), 9.00, (d, 1H), 8.78 (dd, 1H), 8.23 (dd, 1H), 8.06 (m, 2H), 7.70 (d, 1H), 7.64 (m, 2H), 7.58 (m, 1H), 7.54 (m, 1H), 4.09 (s, 3H).	MS 345.3 (M+H) <sup>+</sup>
44.		9.22 (s, 1H), 9.00 (d, 1H), 8.93 (d, 1H), 8.89 (d, 1H), 8.83 (d, 1H), 8.72 (dd, 1H), 8.23 (dd, 1H), 8.09-8.16 (m, 3H), 7.89 (d, 1H), 4.09 (s, 3H).	MS 331.3 (M+H) <sup>+</sup>
45.		8.89 (d, 1H), 8.68 (dd, 1H), 8.34 (d, 1H), 8.11-8.15 (m, 3H), 8.07 (dd, 1H), 7.97-8.00 (m, 2H), 4.12 (s, 3H).	MS 336.8 (M+H) <sup>+</sup>
46.		8.94 (t, 1H), 8.82 (d, 1H), 8.41 (dd, 1H), 8.31 (d, 1H), 8.19 (dd, 1H), 8.18 (d, 1H), 8.16 (d, 1H), 8.14 (s, 1H), 8.09 (dt, 1H), 7.88 (t, 1H), 7.64	MS 340 (M+H) <sup>+</sup>

EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
		(ddd, 1H), 7.29 (dd, 1H), 6.83 (d, 1H)	
47.		8.98 (br s, 1H), 8.83 (d, 1H), 8.45 (dd, 1H), 8.32 (d, 1H), 8.26 (dt, 1H), 8.26 (d, 1H), 8.16 (dd, 1H), 8.10 (dt, 1H), 8.01 (dt, 1H), 7.64 (ddd, 1H), 7.32 (dd, 1H), 6.86 (d, 1H)	MS 358 (M+H) <sup>+</sup>
48.		8.84 (d, 1H), 8.35 (dd, 1H), 8.32 (s, 1H), 8.31 (dd, 1H), 8.23 (dd, 1H), 8.15 (dd, 1H), 8.14 – 8.08 (m, 2H), 7.87 (dd, 1H), 7.64 (ddd, 1H), 7.27 (dd, 1H), 6.83 (d, 1H)	MS 358 (M+H) <sup>+</sup>
49.		8.60 (s, 1H), 8.52-8.51 (d, 1H), 8.364-8.358 (t, 1H), 7.923-7.919 (t, 1H), 7.88-7.786 (m, 1H), 7.77-7.72 (m, 4H), 7.697-7.658 (m, 2H), 7.56-7.53 (m, 2H), 7.50-7.47 (m, 1H).	MS 410.97 (M <sup>+</sup> +H).
50.		8.85 (s, 1H), 8.68-8.67 (d, 1H), 8.33-8.31 (dd, 1H), 8.27-8.55 (d, 1H), 8.17-78.00 (m, 7H), 7.54-7.46 (m, 3H), 7.31-7.30 (m, 1H).	MS 382.41 (M <sup>+</sup> +H).
51.		9.60 (d, 1H), 9.01 (d, 1H), 8.99 (d, 1H), 8.87 (d, 1H), 8.78 (m, 1H), 8.75 (d, 1H), 8.39 (d, 1H), 8.02 (d, 1H), 7.94 (dd, 1H), 7.50 (m, 2H).	MS 302.2 (M <sup>+</sup> +H).
52.		9.51 (d, 1H), 8.90 (d, 1H), 8.85 (d, 1H), 8.70 (dd, 1H), 8.36 (d, 1H), 7.92 (dd, 1H), 7.46 (m, 1H), 7.25 (dd, 1H), 7.19 (d, 1H), 6.91 (dd, 1H), 6.84 (d, 1H), 3.89 (s, 2H).	MS 316.5 (M <sup>+</sup> +H).







EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
53.		9.66 (d, 1H), 8.97 (s, 1H), 8.86 (d, 1H), 8.79 (m, 1H), 8.38 (d, 1H), 7.94 (dd, 1H), 7.87 (d, 1H), 7.78 (dd, 1H), 7.63 (m, 2H), 7.47 (m, 1H).	MS 326.4 (M <sup>+</sup> +H).
54.		9.51 (d, 1H), 8.79 (d, 1H), 8.74 (d, 1H), 8.68 (m, 1H), 8.49 (m, 2H), 8.36 (d, 1H), 8.03 (dd, 1H), 7.58 (m, 1H), 7.40 (d, 1H), 2.42 (s, 3H).	MS 316.0 (M <sup>+</sup> +H).
55.		9.06 (d, 1H), 9.00 (d, 1H), 8.79 (dd, 1H), 8.23 (dd, 1H), 8.09 (m, 2H), 7.70 (d, 1H), 7.64 (m, 2H), 7.58 (m, 1H), 7.53 (m, 1H), 4.07 (s, 3H).	MS 345.3 (M <sup>+</sup> +H).
56.		9.25 (d, 2H), 8.94 (s, broad, 1H), 8.70 (s, broad, 1H), 8.49 (m, 2H), 8.18 (s, 1H), 8.13 (d, 1H), 7.97 (s, broad, 1H), 7.91 (dd, 1H), 4.23 (s, 3H).	MS 331.9 (M <sup>+</sup> +H).
57.		ND	MS 346.3 (M <sup>+</sup> +H).
58.		9.06 (m, 2H), 8.81-8.92 (m, 3H), 8.25 (m, 2H), 8.16 (d, 1H), 8.14 (s, 1H), 7.93 (d, 1H), 4.07 (s, 3H).	MS 374.3 (M <sup>+</sup> +H).
59.		8.86-8.92 (m, 2H), 8.39 (d, 1H), 7.91-8.08 (m, 4H), 7.69 (d, 1H), 7.43-7.51 (m, 2H), 4.05 (s, 3H).	MS 356.4 (M <sup>+</sup> +H).







EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
60.		8.89 (m, 2H), 8.69 (d, 1H), 8.39 (d, 1H), 8.11 (d, 1H), 8.02 (d, 1H), 7.94 (dd, 1H), 7.73 (d, 1H), 7.48 (m, 1H), 4.08 (s, 3H).	MS 357.3 (M <sup>+</sup> +H).
61.		8.90 (d, 1H), 8.39 (d, 1H), 7.89-8.00 (m, 4H), 7.79 (d, 1H), 7.66 (d, 1H), 7.56 (m, 1H), 7.49 (m, 2H), 3.96 (s, 3H).	MS 355.3 (M <sup>+</sup> +H).
62.		8.88 (d, 1H), 8.38 (d, 1H), 8.00 (d, 1H), 7.92 (m, 2H), 7.73 (d, 2H), 7.69 (d, 2H), 7.49 (m, 2H), 3.98 (s, 3H).	MS 355.3 (M <sup>+</sup> +H).
63.		9.02 (d, 1H), 8.37 (d, 1H), 8.21 (d, 1H), 7.99 (d, 1H), 7.91 (m, 2H), 7.63 (m, 2H), 7.46 (m, 1H), 6.72 (m, 1H), 4.04 (s, 3H), 4.03 (s, 3H).	MS 361.1 (M <sup>+</sup> +H).
64.		8.95 (d, 1H), 8.88 (d, 1H), 8.68 (m, 1H), 8.41 (d, 1H), 8.33 (s, 1H), 8.20 (s, 1H), 8.01 (m, 1H), 7.95 (m, 1H), 7.56 (s, 1H), 7.45 (m, 2H), 2.59 (s, 3H).	MS 315.4 (M <sup>+</sup> +H).
65.		8.88 (d, 1H), 8.78 (d, 1H), 8.41 (d, 1H), 8.32 (s, 1H), 7.96 (m, 2H), 7.82 (m, 1H), 7.50 (m, 1H), 7.34 (m, 2H), 3.95 (s, 3H).	MS 411.0 (M <sup>+</sup> +H+2)
66.		8.92-8.94 (m, 1H), 8.77-8.79 (m, 1H), 8.38-8.40 (m, 1H), 8.11-8.16 (m, 3H), 8.02-8.04 (m, 1H), 7.62 (s, 1H)	MS 334.3 (M <sup>+</sup> +H).
67.		8.8 (d, 1H), 8.18-8.20 (m, 1H), 8.09-8.10 (m, 1H), 8.05 (d, 1H), 7.97 (d, 1H), 7.92 (m, 1H), 7.90 (s, 1H)	MS 334.2 (M <sup>+</sup> +H).







EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
68.		8.81-8.91 (m, 1H), 8.82-8.83 (m, 1H), 8.69-8.70 (d, 1H), 8.56-8.58 (m, 1H), 8.5-8.52 (d, 1H), 8.45-8.48 (dd, 1H), 8.22-8.4 (m, 3H), 8.29 (ddd, 1H), 8.20 (m, 1H), 7.88-8.03 (m, 2H), 7.78-7.80 (m, 1H)	MS 394.3 (M <sup>+</sup> +H).
69.		9.18 (m, 1H), 8.6 (s, 1H), 8.49 (ddd, 1H), 8.38 (m, 2H), 8.22 (m, 1H), 7.87 (m, 2H), 7.57 (m, 2H), 7.50 (m, 1H)	MS 343 (M <sup>+</sup> +H).
70.		8.82-8.83 (m, 1H), 8.79 (m, 1H), 8.25-8.34 (m, 5H), 8.08 (ddd, 1H), 7.92 (dd, 1H), 7.64 (dd, 1H)	MS 291 (M <sup>+</sup> +H).
71.		8.88 (s, 1H), 8.81-8.79 (d, 1H), 8.71-8.68 (m, 1H), 8.46-8.45 (m, 1H), 8.34-8.28 (m, 2H), 8.24-8.19 (m, 1H), 8.14-8.11 (d, 1H), 7.83-7.79 (m, 1H), 7.71-7.67 (m, 1H), 2.67 (s, 3H)	MS 349.1 (M <sup>+</sup> +H).
72.		8.76-8.72 (m, 1H), 8.45-8.42 (m, 1H), 8.41-8.38 (m, 1H), 8.28-8.25 (d, 1H), 8.07-8.02 (m, 1H), 7.84-7.81 (m, 1H), 7.73-7.68 (m, 2H), 7.61-7.59 (d, 1H), 7.56-7.51 (m, 2H)	MS 359.0 (M <sup>+</sup> +H).
73.		9.39 (s, 1H), 8.80-8.81 (d, 1H), 8.73-8.74 (d, 1H), 8.57 (s, 1H), 8.44-8.45 (d, 1H), 8.35-8.37 (d, 1H), 8.25-8.30 (m, 2H), 8.04-8.09 (m, 2H), 7.97-8.00 (dd, 1H), 7.61-7.63 (dd, 1H), 7.26 (d, 1H).	MS 340 (M <sup>+</sup> +H).







EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
74.		9.50 (s, 1H), 8.82-8.83 (m, 1H), 8.75-8.76 (d, 1H), 8.48 (m, 2H), 8.27-8.34 (m, 3H), 8.09-8.14 (m, 2H), 7.64-7.66 (dd, 1H), 7.29-7.30 (d, 1H).	MS 359 (M <sup>+</sup> +H)
75.		ND	MS 341 (M <sup>+</sup> +H)
76.		ND	MS 370 (M <sup>+</sup> +H)
77.		8.92 – 8.93 (m, 1H), 8.80 – 8.82 (m, 1H), 8.75 – 8.77 (m, 1H), 8.23 – 8.34 (m, 3H), 8.14 – 8.17 (m, 1H), 8.07 (ddd, 1H, J = 10.0, 10.0, 2.4 Hz), 7.97 (ddd, 1H, J = 10.4, 10.4, 2.4 Hz), 7.82 (dd, 1H, J = 10.8, 10.8 Hz), 7.62 (ddd, 1H, J = 10.0, 6.4, 1.2), 7.44 – 7.48 (m, 1H).	MS 301 (M <sup>+</sup> +H)
78.		9.40-9.42 (d, 1H), 9.15-9.16 (d, 1H), 8.92-8.93 (d, 1H), 8.67-8.68 (d, 1H), 8.57-8.60 (m, 1H), 8.49-8.54 (m, 2H), 8.27-8.29 (d, 1H), 8.21-8.24 (dd, 1H), 8.15-8.18 (1H), 8.01-8.04 (t, 1H), 7.93-8.00 (m, 2H).	MS 351 (M <sup>+</sup> +H)
79.		8.99 (m, 1H), 8.81-8.83 (d, 1H), 8.69-8.72 (m, 1H), 8.37-8.39 (dd, 1H), 8.27 (s, 1H), 8.13-8.15 (m, 1H), 7.89-7.91 (d, 1H), 7.74-7.89 (m, 2H), 7.64-7.70 (m, 2H), 7.52-7.53 (d, 1H).	MS 368 (M <sup>+</sup> +H)









EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
80.		8.82-8.83 (d, 1H), 8.30-8.31 (d, 1H), 8.19-8.21 (m, 2H), 8.06-8.08 (m, 3H), 7.79-7.83 (m, 1H), 7.67-7.68 (d, 1H), 7.58-7.61 (m, 1H), 6.70-6.71 (d, 1H), 6.02 (s, 2H).	MS 316 (M <sup>+</sup> +H)
81.		8.87-8.88 (m, 1H), 8.38-8.41 (m, 2H), 8.32-8.33 (m, 1H), 8.16-8.17 (dd, 1H), 7.93-7.95 (m, 1H), 7.68-7.71 (m, 2H), 7.47-7.50 (m, 1H), 6.82-6.85 (dd, 1H), 4.64 (s, 2H).	MS 316 (M <sup>+</sup> +H)
82.		8.95-8.97 (m, 1H), 8.75-8.78 (m, 1H), 8.60-8.63 (m, 1H), 8.52 (m, 1H), 8.43 (m, 1H), 8.07 (m, 1H), 7.95-7.97 (d, 1H), 7.88-7.90 (m, 2H), 7.83-7.86 (m, 1H), 7.76-7.77 (d, 1H), 7.66-7.96 (dd, 1H).	MS 325 (M <sup>+</sup> +H)
83.		8.85-8.86 (d, 1H), 8.33-8.34 (d, 1H), 8.26-8.28 (m, 1H), 8.23-8.26 (m, 1H), 8.10-8.14 m, 1H), 8.08-8.10 (d, 1H), 7.85-7.92 (m, 3H), 7.74-7.76 (dd, 1H), 7.66-7.68 (m, 1H).	MS 344 (M <sup>+</sup> +H)
84.		9.38 (s, 1H), 8.92-8.93 (d, 1H), 8.86-8.88 (m, 1H), 8.83-8.84 (d, 1H), 8.52 (s, 1H), 8.32-8.33 (d, 1H), 8.21-8.23 (m, 1H), 8.09-8.13 (m, 2H), 8.03-8.06 (m, 1H), 7.64-7.66 (m, 1H).	MS 319 (M <sup>+</sup> +H)
85.		8.82-8.83 (m, 1H), 8.30-8.32 (d, 1H), 8.09-8.14 (m, 3H), 7.78-7.80 (d, 1H), 7.61-7.65 (m, 3H), 7.53-7.57 (m, 3H), 7.40-7.42 (m, 1H).	MS 333 (M <sup>+</sup> +H)




EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
86.		8.86-8.88 (m, 1H), 8.51-8.53 (dd, 1H). 8.38-8.39 (d, 1H), 8.22 (s, 1H), 8.12-8.14 (m, 1H), 7.92-7.96 (m, 1H), 7.78-7.81 (dd, 1H), 7.47-7.50 (m, 1H), 7.39-7.43 (m, 2H).	MS 354 (M <sup>+</sup> +H)
87.		8.87-8.88 (d, 1H), 8.35-8.40 (m, 3H), 7.92-7.95 (m, 1H), 7.67-7.70 (m, 2H), 7.58-7.60 (d, 1H), 7.46-7.49 (dd, 1H), 7.16-7.19 (dd, 1H), 7.02-7.06 (m, 1H).	MS 396 (M <sup>+</sup> +H)
88.		8.87-8.88 (m, 1H), 8.39-8.40 (d, 1H), 8.24-8.25 (d, 1H), 8.09 (s, 1H), 7.93-7.94 (m, 1H), 7.82-7.83 (m, 1H), 7.73-7.75 (dd, 1H), 7.44-7.48 (m, 1H), 7.30 (s, 1H), 7.03-7.05 (dd, 1H).	MS 361 (M <sup>+</sup> +H)
89.		8.82-8.83 (m, 1H), 8.42 (s, 1H), 8.36-8.37 (m, 1H), 8.29-8.31 (m, 1H), 8.15-8.17 (m, 1H), 8.08-8.09 (m, 1H), 7.90-7.91 (m, 2H), 7.75-7.78 (m, 1H), 7.57-7.58 (m, 1H), 7.34-7.35 (m, 1H).	MS 343 (M <sup>+</sup> +H)
90.		8.89-8.91 (m, 1H), 8.60-8.62 (m, 1H), 8.51 (s, 1H), 8.37-8.41 (m, 2H), 8.11-8.12 (d, 1H), 7.93-7.96 (d, 1H), 7.88-7.90 (m, 1H), 7.80-7.83 (m, 2H).	MS 334 (M <sup>+</sup> +H)
91.		8.82-8.83 (d, 1H), 8.72-8.73 (d, 1H), 8.33 (s, 1H), 8.30-8.32 (d, 1H), 8.24-8.26 (d, 1H), 8.21-8.23 (d, 1H), 8.08-8.11 (m, 1H), 7.86-7.88 (d, 1H), 7.73-7.76 (dd, 1H), 7.63-7.65 (dd, 1H), 2.51 (s, 3H).	MS 333 (M <sup>+</sup> +H)




EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
92.		8.92-8.93 (d, 1H), 8.62-8.64 (d, 1H), 8.39-8.43 (m, 2H), 7.89-7.96 (m, 2H), 7.77-7.79 (m, 1H), 7.65-7.67 (d, 1H), 7.56-7.60 (m, 1H).	MS 361 (M <sup>+</sup> +H)
93.		13.84 (m, 1H), 8.82-8.83 (d, 1H), 8.30-8.31 (d, 1H), 8.19-8.21 (d, 1H), 8.17 (s, 1H), 8.12-8.14 (d, 1H), 8.08-8.10 (m, 1H), 8.04-8.06 (m, 2H), 7.69-7.71 (d, 1H), 7.63-7.65 (m, 1H), 7.04-7.07 (dd, 1H).	MS 334 (M <sup>+</sup> +H)
94.		13.84 (m, 1H), 8.82 (s, 1H), 8.53 (m, 1H), 8.30-8.34 (m, 2H), 8.20-8.21 (m, 2H), 8.08-8.13 (m, 2H), 7.63-7.69 (m, 2H), 7.57 (m, 1H), 7.03-7.05 (d, 1H).	MS 334 (M <sup>+</sup> +H)
95.		9.02 (s, 1H), 8.92-8.93 (d, 1H), 8.81-8.82 (d, 1H), 8.26-8.30 (m, 2H), 8.20 (m, 1H), 8.08-8.11 (m, 2H), 7.81-7.83 (dd, 1H), 7.63-7.65 (dd, 1H), 2.48 (s, 3H).	MS 333 (M <sup>+</sup> +H)
96.		8.94 (s, 1H), 8.86-8.87 (d, 1H), 8.82-8.83 (d, 1H), 8.30-8.31 (d, 1H), 8.24-8.26 (m, 2H), 8.08-8.11 (m, 1H), 8.05-8.07 (d, 1H), 7.80-7.82 (m, 1H), 7.63-7.66 (dd, 1H), 2.58 (s, 3H).	MS 333 (M <sup>+</sup> +H)
97.		9.00-9.02 (m, 1H), 8.83-8.84 (m, 2H), 8.70-8.73 (m, 1H), 8.46 (s, 1H), 8.29-8.31 (d, 1H), 8.23-8.24 (d, 1H), 8.14-8.16 (m, 1H), 7.85-7.88 (dd, 1H), 7.79-7.81 (d, 1H), 7.64-7.68 (m, 1H), 7.56-7.59 (m, 1H).	MS 344 (M <sup>+</sup> +H)

EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
98.		9.38 (s, 1H), 8.96-8.97 (d, 1H), 8.86-8.88 (d, 1H), 8.55 (s, 1H), 8.19-8.22 (m, 2H), 8.07-8.09 (dd, 1H), 7.86 (s, 2H).	MS 308 (M <sup>+</sup> +H)
99.		8.87-8.88 (d, 1H), 8.81-8.82 (d, 1H), 8.55-8.57 (d, 1H), 8.29-8.30 (d, 1H), 8.23-8.25 (m, 2H), 8.08-8.11 (m, 1H), 8.00-8.03 (m, 1H), 7.78-7.80 (d, 1H), 7.63-7.65 (dd, 1H), 2.77 (s, 3H).	MS 333 (M <sup>+</sup> +H)
100.		8.88 (s, 1H), 8.83-8.84 (d, 1H), 8.16-8.19 (m, 2H), 7.99-8.01 (d, 1H), 7.82-7.84 (d, 1H), 7.65 (s, 2H), 2.55 (s, 3H).	MS 322 (M <sup>+</sup> +H)
101.		9.19-9.21 (m, 1H), 8.79-8.81 (m, 2H), 8.76 (s, 1H), 8.50 (m, 1H), 8.29-8.31 (d, 1H), 8.18-8.19 (d, 1H), 8.07-8.09 (m, 2H), 7.58-7.62 (m, 3H), 7.54-7.56 (m, 1H), 2.54 (s, 3H).	MS 333 (M <sup>+</sup> +H)
102.		9.29 (s, 1H), 8.90-8.91 (d, 1H), 8.79-8.81 (m, 1H), 8.51 (s, 1H), 8.29-8.31 (d, 1H), 8.18-8.20 (d, 1H), 8.09-8.11 (m, 2H), 8.01-8.03 (d, 1H), 7.53-7.62 (m, 4H), 2.80 (s, 3H).	MS 333 (M <sup>+</sup> +H)
103.		8.82-8.83 (d, 1H), 8.78 (s, 1H), 8.65 (m, 1H), 8.31-8.33 (d, 1H), 8.19-8.22 (m, 2H), 8.08-8.14 (m, 2H), 7.92-7.94 (dd, 1H), 7.60-7.66 (m, 1H).	MS 333 (M <sup>+</sup> +H)





EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
104.		9.41-9.42 (d, 1H), 8.71 (d, 1H), 8.36 (s, 2H), 8.24 (s, 2H), 7.87 (s, 1H), 7.09-7.10 (d, 1H).	MS 323 (M <sup>+</sup> +H)
105.		8.34 (s, 1H), 8.24-8.26(d, 1H), 8.17-8.19 (d, 1H), 8.14 (s, 1H), 7.85 (s, 2H), 7.75-7.76 (m, 1H), 7.65 (s, 1H), 7.18-7.20 (d, 1H).	MS 356 (M <sup>+</sup> +H)
106.		8.92-8.97 (m, 1H), 8.79-8.80 (d, 1H), 8.64-8.67 (dd, 1H), 8.09-8.11 (m, 1H), 7.90-7.92 (m, 2H), 7.47-7.49 (d, 1H), 7.38-7.41 (m, 1H), 7.23-7.25 (m, 1H), 7.09-7.10 (d, 1H), 7.02-7.05 (m, 1H), 3.94 (s, 3H), 3.80 (s, 3H).	MS 360 (M <sup>+</sup> +H)
107.		8.98-9.00 (m, 1H), 8.72-8.74 (d, 1H), 8.53-8.56 (m, 1H), 7.98-8.05 (m, 3H), 7.86-7.87 (d, 1H), 7.78-7.82 (m, 1H), 7.56-7.63 (m, 3H), 4.03 (s, 3H).	MS 355 (M <sup>+</sup> +H)
108.		ND	MS 335.2 (M <sup>+</sup> +H).
109.		8.73-8.71 (m, 2H), 8.31-8.2 (m, 3H), 8.0 (dt, 1H), 7.78 (dt, 1H), 7.58 (d, 1H), 7.5-7.43 (m, 2H), 7.3-7.27 (m, 1H), 2.5 (s, 3H).	MS 315.4 (M <sup>+</sup> +H).





EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
110.		8.7 (br s, 2H), 8.28 (d, 1H), 8.10-8.0 (m, 2H), 7.9-7.8 (m, 2H), 7.8 (d, 1H), 7.75 (t, 1H), 7.52-7.48 (m, 1H), 7.3-7.2 (m, 1H), 4.1 (s, 3H).	MS 331.3 (M <sup>+</sup> +H).
111.		ND	MS 331.8 (M <sup>+</sup> +H).
112.		9.14-9.13 (d, 1H) 8.79-8.76 (d, 1H), 8.56-8.52 (m, 3H), 8.34-8.33 (d, 1H), 8.22-8.20 (d, 1H), 8.04-8.01 (t, 1H), 7.69-7.68 (d, 1H) 7.60-7.57 (m, 2H), 7.02-7.01 (d, 1H), 4.14 (s, 3H).	MS 370.1 (M <sup>+</sup> +H).
113.		9.00-8.99 (d, 1H), 8.84-8.83 (d, 1H), 8.75-8.72 (t, 1H), 8.28-8.27 (d, 1H), 8.17-8.15 9t, 1H), 8.09 (s, 1H), 8.02-7.96 (m, 2H), 7.79 (s, 1H), 6.57-6.56 (t, 1H), 4.10 (s, 3H).	MS 320.3 (M <sup>+</sup> +H).
114.		9.04-9.07(m, 2H), 8.82-8.87 (m, 2H), 8.74-8.77 (m, 1H), 8.45-8.48 (m, 3H), 8.17-8.26 (m, 3H).	MS 319.9 (M <sup>+</sup> +H)
115.		8.81-8.83 (m, 2H), 8.73-8.74 (d, 1H), 8.58-8.61 (dd, J=6, 6, 1H), 8.31-8.33, J=6, 1H), 8.23-8.25 (d, J=6. 1H), 8.07-8.09 (m, 2H), 8.02-8.00 (d, J=6, 1H), 7.63-7.65 (t, 1H), 7.57-7.59 (t, 1H).	MS 326.2 (M <sup>+</sup> +H)
116.		8.85 (s, 1H), 8.76-8.77 (d, J=3, 1H), 8.56-8.60 (m, 2H), 8.33-8.35 (d, J=6, 1H), 8.05-8.12 (m, 2H), 7.94-7.96 (d, J=6, 1H), 7.69-7.71 (d, J=6,	MS 369.2 (M <sup>+</sup> +H)





EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
		1H), 7.64-7.66 (m, 1H), 7.58-7.60 (m, 1H).	
117.		8.86-8.87 (d, J=3, 1H), 8.77-8.78 (d, J=3, 1H), 8.38-8.40 (d, J=6, 1H), 8.34 (s, 1H), 8.12-8.14 (d, J=6, 1H), 7.92-7.94 (t, 1H), 7.81-7.84 (m, 1H), 7.77-7.79 (d, J=6, 1H), 7.48-7.50 (t, 1H), 7.35-7.37 (t, 1H).	MS 385.1 (M <sup>+</sup> +H)
118.		ND	MS 346 (M+H)
119.		ND	MS 370 (M+H)





EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
120.		ND	MS 370 (M+H)
121.		ND	MS 385 (M+H)
122.		ND	MS 369 (M+H)













EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
123.		ND	MS 331 (M+H)
124.		ND	MS 315 (M+H)
125.		ND	MS 315 (M+H)
126.		ND	MS 343 (M+H)




EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
127.		ND	MS 319 (M+H)
128.		ND	MS 337 (M+H)
129.		ND	MS 316 (M+H)
130.		ND	MS 317 (M+H)





EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
131.		ND	MS 319 (M+H)
132.		ND	MS 326 (M+H)
133.		ND	MS 329 (M+H)
134.		ND	MS 329 (M+H)

EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
135.		ND	MS 329 (M+H)
136.		ND	MS 331 (M+H)
137.		ND	MS 337 (M+H)
138.		ND	MS 344 (M+H)




EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
139.		ND	MS 314 (M <sup>+</sup> +H)
140.		ND	MS 318 (M <sup>+</sup> +H)
141.		ND	MS 318 (M <sup>+</sup> +H)
142.		ND	MS 332 (M <sup>+</sup> +H)
143.		ND	MS 336 (M <sup>+</sup> +H)
144.		ND	MS 376 (M <sup>+</sup> +H)





EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
145.		ND	MS 319 (M <sup>+</sup> +H)
146.		ND	MS 332 (M <sup>+</sup> +H)
147.		ND	MS 387 (M <sup>+</sup> +H)
148.		ND	MS 343 (M <sup>+</sup> +H)




EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
149.		ND	MS 354 (M <sup>+</sup> +H)
150.		ND	MS 352 (M <sup>+</sup> +H)
151.		ND	MS 378 (M <sup>+</sup> +H)





EXAMPLE	Structure	$^1\text{H}$ NMR	MS (ESI)
152.		ND	MS 346 ( $\text{M}^+ + \text{H}$ )
153.		ND	MS 352 ( $\text{M}^+ + \text{H}$ )
154.		ND	MS 360 ( $\text{M}^+ + \text{H}$ )
155.		ND	MS 346 ( $\text{M}^+ + \text{H}$ )











EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
156.		ND	MS 346 (M <sup>+</sup> +H)
157.		ND	MS 352 (M <sup>+</sup> +H)
158.		ND	MS 334 (M <sup>+</sup> +H)





EXAMPLE	Structure	$^1\text{H NMR}$	MS (ESI)
159.		ND	MS 346 ( $\text{M}^+ + \text{H}$ )
160.		ND	MS 348 ( $\text{M}^+ + \text{H}$ )
161.		ND	MS 336 ( $\text{M}^+ + \text{H}$ )
162.		ND	MS 361 ( $\text{M}^+ + \text{H}$ )





EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
163.		ND	MS 363 (M <sup>+</sup> +H)
164.		ND	MS 387 (M <sup>+</sup> +H)
165.		ND	MS 348 (M <sup>+</sup> +H)

EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
166.		ND	MS 361 (M <sup>+</sup> +H)
167.		ND	MS 354 (M <sup>+</sup> +H)
168.		ND	MS 336 (M <sup>+</sup> +H)
169.		ND	MS 362 (M <sup>+</sup> +H)





EXAMPLE	Structure	$^1\text{H NMR}$	MS (ESI)
170.		ND	MS 334 ( $\text{M}^+ + \text{H}$ )
171.		ND	MS 328 ( $\text{M}^+ + \text{H}$ )
172.		ND	MS 345 ( $\text{M}^+ + \text{H}$ )
173.		ND	MS 314 ( $\text{M}^+ + \text{H}$ )




EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
174.		ND	MS 314 (M <sup>+</sup> +H)
175.		ND	MS 342 (M <sup>+</sup> +H)
176.		ND	MS 334 (M <sup>+</sup> +H)
177.		ND	MS 318 (M <sup>+</sup> +H)





EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
178.		ND	MS 369 (M <sup>+</sup> +H)
179.		ND	MS 336 (M <sup>+</sup> +H)
180.		ND	MS 369 (M <sup>+</sup> +H)
181.		ND	MS 316 (M <sup>+</sup> +H)




EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
182.		ND	MS 328 (M <sup>+</sup> +H)
183.		ND	MS 328 (M <sup>+</sup> +H)
184.		ND	MS 328 (M <sup>+</sup> +H)
185.		ND	MS 336 (M <sup>+</sup> +H)











EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
186.		ND	MS 342 (M <sup>+</sup> +H)
187.		ND	MS 343 (M <sup>+</sup> +H)
188.		ND	MS 330 (M <sup>+</sup> +H)
189.		ND	MS 316 (M <sup>+</sup> +H)




EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
190.		ND	MS 318 (M <sup>+</sup> +H)
191.		ND	MS 325 (M <sup>+</sup> +H)
192.		ND	MS 330 (M <sup>+</sup> +H)




EXAMPLE	Structure	$^1\text{H}$ NMR	MS (ESI)
193.		ND	MS 362 ( $\text{M}^+\text{+H}$ )
194.		ND	MS 362 ( $\text{M}^+\text{+H}$ )
195.		ND	MS 364 ( $\text{M}^+\text{+H}$ )
196.		ND	MS 375 ( $\text{M}^+\text{+H}$ )

EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
197.		ND	MS 376 (M <sup>+</sup> +H)
198.		ND	MS 408 (M <sup>+</sup> +H)
199.		ND	MS 410 (M <sup>+</sup> +H)




EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
200.		ND	MS 386 (M <sup>+</sup> +H)
201.		ND	MS 320 (M <sup>+</sup> +H)
202.		ND	MS 350 (M <sup>+</sup> +H)
203.		ND	MS 384 (M <sup>+</sup> +H)





EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
204.		ND	MS 384 (M <sup>+</sup> +H)
205.		ND	MS 354 (M <sup>+</sup> +H)
206.		ND	MS 348 (M <sup>+</sup> +H)
207.		ND	MS 344 (M <sup>+</sup> +H)








EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
208.		ND	MS 346 (M <sup>+</sup> +H)
209.		ND	MS 357 (M <sup>+</sup> +H)
210.		ND	MS 357 (M <sup>+</sup> +H)

EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
211.		ND	MS 358 (M <sup>+</sup> +H)
212.		ND	MS 369 (M <sup>+</sup> +H)
213.		ND	MS 384 (M <sup>+</sup> +H)

















EXAMPLE	Structure	$^1\text{H}$ NMR	MS (ESI)
214.		ND	MS 392 ( $\text{M}^+ + \text{H}$ )
215.		ND	MS 393 ( $\text{M}^+ + \text{H}$ )
216.		ND	MS 363 ( $\text{M}^+ + \text{H}$ )




EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
217.		ND	MS 345 (M <sup>+</sup> +H)
218.		ND	MS 351 (M <sup>+</sup> +H)
219.		ND	MS 332 (M <sup>+</sup> +H)
220.		ND	MS 368 (M <sup>+</sup> +H)

EXAMPLE	Structure	$^1\text{H}$ NMR	MS (ESI)
221.		ND	MS 331 ( $\text{M}^+ + \text{H}$ )
222.		ND	MS 344 ( $\text{M}^+ + \text{H}$ )
223.		ND	MS 348 ( $\text{M}^+ + \text{H}$ )
224.		ND	MS 360 ( $\text{M}^+ + \text{H}$ )
225.		ND	MS 388 ( $\text{M}^+ + \text{H}$ )
226.		ND	MS 331 ( $\text{M}^+ + \text{H}$ )
227.		ND	MS 364 ( $\text{M}^+ + \text{H}$ )

EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
228.		ND	MS 358 (M <sup>+</sup> +H)
229.		ND	MS 360 (M <sup>+</sup> +H)
230.		ND	MS 344 (M <sup>+</sup> +H)
231.		ND	MS 380 (M <sup>+</sup> +H)
232.		ND	MS 372 (M <sup>+</sup> +H)
233.		ND	MS 348 (M <sup>+</sup> +H)
234.		ND	MS 366 (M <sup>+</sup> +H)


EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
235.		ND	MS 346 (M <sup>+</sup> +H)
236.		ND	MS 348 (M <sup>+</sup> +H)
237.		ND	MS 358 (M <sup>+</sup> +H)
238.		ND	MS 366 (M <sup>+</sup> +H)
239.		ND	MS 372 (M <sup>+</sup> +H)
240.		ND	MS 373 (M <sup>+</sup> +H)
241.		ND	MS 360 (M <sup>+</sup> +H)





EXAMPLE	Structure	$^1\text{H}$ NMR	MS (ESI)
242.		ND	MS 331 ( $\text{M}^+ + \text{H}$ )
243.		ND	MS 374 ( $\text{M}^+ + \text{H}$ )
244.		ND	MS 381 ( $\text{M}^+ + \text{H}$ )
245.		ND	MS 387 ( $\text{M}^+ + \text{H}$ )
246.		ND	MS 399 ( $\text{M}^+ + \text{H}$ )
247.		ND	MS 399 ( $\text{M}^+ + \text{H}$ )
248.		ND	MS 332 ( $\text{M}^+ + \text{H}$ )

EXAMPLE	Structure	$^1\text{H}$ NMR	MS (ESI)
249.		ND	MS 396 ( $\text{M}^+ + \text{H}$ )
250.		ND	MS 398 ( $\text{M}^+ + \text{H}$ )
251.		ND	MS 360 ( $\text{M}^+ + \text{H}$ )





Examples 252-281 have mGluR5 inhibitory activity greater than 1  $\mu\text{M}$  in the calcium flux assay:




5




EXAMPLE	Structure	$^1\text{H}$ NMR	MS (ESI)
252.		ND	MS 345 ( $\text{M} + \text{H}$ )




EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
257.		ND	MS 358 (M+H)
258.		ND	MS 358 (M+H)
259.		ND	MS 358 (M+H)
260.		ND	MS 359 (M+H)












EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
253.		ND	MS 345 (M+H)
254.		ND	MS 345 (M+H)
255.		ND	MS 347 (M+H)
256.		ND	MS 352 (M+H)




EXAMPLE	Structure	$^1\text{H}$ NMR	MS (ESI)
261.		ND	MS 370 (M+H)
262.		ND	MS 377 (M+H)
263.		ND	MS 391 (M+H)

EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
264.		ND	MS 391 (M+H)
265.		ND	MS 393 (M+H)
266.		ND	MS 394 (M+H)


EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
267.		ND	MS 407 (M+H)
268.		ND	MS 439 (M+H)
269.		ND	MS 369 (M+H)





EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
274.		ND	MS 385 (M+H)
275.		ND	MS 332 (M+H)
276.		ND	MS 370 (M+H)
277.		ND	MS 361 (M+H)
278.		ND	MS 315 (M+H)

EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
270.		ND	MS 357 (M+H)
271.		ND	MS 367 (M+H)
272.		ND	MS 367 (M+H)
273.		ND	MS 359 (M+H)





EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
279.		ND	MS 329 (M+H)
280.		ND	MS 344 (M+H)
281.		ND	MS 345 (M+H)




Examples 282-347 have mGluR5 inhibitory activity greater than 3  $\mu$ M in the calcium flux assay:





EXAMPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
282.		ND	MS 330 (M+H)




EXA MPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
283.		ND	MS 358 (M+H)
284.		ND	MS 348 (M+H)
285.		ND	MS 332 (M+H)
286.		ND	MS 368 (M+H)









EXA MPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
287.		ND	MS 378 (M+H)
288.		ND	MS 334 (M+H)
289.		ND	MS 346 (M+H)
290.		ND	MS 334 (M+H)





EXA MPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
291.		ND	MS 350 (M+H)
292.		ND	MS 360 (M+H)
293.		ND	MS 360 (M+H)




EXA MPLE	Structure	$^1\text{H}$ NMR	MS (ESI)
294.		ND	MS 328 (M+H)
295.		ND	MS 343 (M+H)
296.		ND	MS 344 (M+H)
297.		ND	MS 362 (M+H)

EXA MPLE	Structure	$^1\text{H}$ NMR	MS (ESI)
298.		ND	MS 387 (M+H)
299.		ND	MS 387 (M+H)
300.		ND	MS 394 (M+H)






EXA MPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
301.		ND	MS 402 (M+H)
302.		ND	MS 408 (M+H)
303.		ND	MS 411 (M+H)








EXA MPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
304.		ND	MS 424 (M+H)
305.		ND	MS 386 (M+H)
306.		ND	MS 376 (M+H)








EXA MPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
307.		ND	MS 402 (M+H)
308.		ND	MS 349 (M+H)
309.		ND	MS 344 (M+H)
310.		ND	MS 344 (M+H)







EXA MPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
311.		ND	MS 357 (M+H)
312.		ND	MS 369 (M+H)
313.		ND	MS 376 (M+H)












EXA MPLE	Structure	$^1\text{H}$ NMR	MS (ESI)
314.		ND	MS 390 (M+H)
315.		ND	MS 390 (M+H)
316.		ND	MS 364 (M+H)
317.		ND	MS 375 (M+H)
318.		ND	MS 344 (M+H)

EXA MPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
319.		ND	MS 364 (M+H)
320.		ND	MS 399 (M+H)
321.		ND	MS 390 (M+H)
322.		ND	MS 390 (M+H)
323.		ND	MS 346 (M+H)
324.		ND	MS 358 (M+H)
325.		ND	MS 358 (M+H)

EXA MPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
326.		ND	MS 358 (M+H)
327.		ND	MS 374 (M+H)
328.		ND	MS 387 (M+H)
329.		ND	MS 387 (M+H)
330.		ND	MS 388 (M+H)
331.		ND	MS 399 (M+H)
332.		ND	MS 406 (M+H)

EXA MPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
333.		ND	MS 414 (M+H)
334.		ND	MS 420 (M+H)
335.		ND	MS 420 (M+H)
336.		ND	MS 422 (M+H)
337.		ND	MS 423 (M+H)
338.		ND	MS 436 (M+H)


EXA MPLE	Structure	<sup>1</sup> H NMR	MS (ESI)
339.		ND	MS 468 (M+H)
340.		ND	MS 398 (M+H)
341.		ND	MS 396 (M+H)
342.		ND	MS 388 (M+H)
343.		ND	MS 414 (M+H)
344.		ND	MS 361 (M+H)
345.		8.91 -8.92(d, 2H), 8.75-8.79 (m, 2H), 8.27-8.29 (d, 2H), 7.97-8.06 (m, 2H), 7.58-7.60 (dd, 1H), 7.46-7.49 (m, 2H).	MS 378 (M <sup>+</sup> +H)

EXA MPLE	Structure	$^1\text{H}$ NMR	MS (ESI)
346.		8.2 (s, 1H), 8.98-9.00 (d, J=6, 1H), 8.84-8.85 (d, J=3, 1H), 8.55-8.59 (m, 2H), 8.33-8.35 (d, J=6, 1H), 8.10-8.12 (t, 1H), 8.64-8.66 (t, 1H), 7.58 (s, 2H), 7.39 (s, 1H), 7.33-7.38 (m, 2H).	MS 340.1 ( $\text{M}^+ + \text{H}$ )
347.		8.85 (br s, 1H), 8.74 (d, 1H), 8.6 (d, 1H), 8.3 (d, 1H), 8.1-8.05 (m, 2H), 7.95 (s, 1H), 7.92 (d, 1H), 7.6-7.55 (m, 1H), 7.5 (d, 1H), 7.4-7.37 (m, 1H), 3.98 (s, 3H).	MS 331.8 ( $\text{M}^+ + \text{H}$ ).

Other variations or modifications, which will be obvious to those skilled in the art, are within the scope and teachings of this invention. This invention is not to be limited except as set forth in the following claims.

## WHAT IS CLAIMED IS:

1. A compound represented by Formula (I):



(I)

or a pharmaceutically acceptable salt thereof, wherein

X and Y each independently is aryl or heteroaryl wherein at least one of X and Y is a heteroaryl with N adjacent to the position of attachment to A or B respectively;

X is optionally substituted with 1-7 independent halogen,  $-\text{CN}$ ,  $\text{NO}_2$ ,  $-\text{C}_1\text{-6alkyl}$ ,  $-\text{C}_1\text{-6alkenyl}$ ,  $-\text{C}_1\text{-6alkynyl}$ ,  $-\text{OR}^1$ ,  $-\text{NR}^1\text{R}^2$ ,  $-\text{C}(=\text{NR}^1)\text{NR}^2\text{R}^3$ ,  $-\text{N}(=\text{NR}^1)\text{NR}^2\text{R}^3$ ,  $-\text{NR}^1\text{COR}^2$ ,  $-\text{NR}^1\text{CO}_2\text{R}^2$ ,  $-\text{NR}^1\text{SO}_2\text{R}^4$ ,  $-\text{NR}^1\text{CONR}^2\text{R}^3$ ,  $-\text{SR}^4$ ,  $-\text{SOR}^4$ ,  $-\text{SO}_2\text{R}^4$ ,  $-\text{SO}_2\text{NR}^1\text{R}^2$ ,  $-\text{COR}^1$ ,  $-\text{CO}_2\text{R}^1$ ,  $-\text{CONR}^1\text{R}^2$ ,  $-\text{C}(=\text{NR}^1)\text{R}^2$ , or  $-\text{C}(=\text{NOR}^1)\text{R}^2$  substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the  $-\text{C}_1\text{-6alkyl}$  substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen,  $-\text{CN}$ ,  $-\text{C}_1\text{-6alkyl}$ ,  $-\text{O}(\text{C}_0\text{-6alkyl})$ ,  $-\text{O}(\text{C}_3\text{-7cycloalkyl})$ ,  $-\text{O}(\text{aryl})$ ,  $-\text{N}(\text{C}_0\text{-6alkyl})(\text{C}_0\text{-6alkyl})$ ,  $-\text{N}(\text{C}_0\text{-6alkyl})(\text{C}_3\text{-7cycloalkyl})$ , or  $-\text{N}(\text{C}_0\text{-6alkyl})(\text{aryl})$  groups;

$\text{R}^1$ ,  $\text{R}^2$ , and  $\text{R}^3$  each independently is  $-\text{C}_0\text{-6alkyl}$ ,  $-\text{C}_3\text{-7cycloalkyl}$ , heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent halogen,  $-\text{CN}$ ,  $-\text{C}_1\text{-6alkyl}$ ,  $-\text{O}(\text{C}_0\text{-6alkyl})$ ,  $-\text{O}(\text{C}_3\text{-7cycloalkyl})$ ,  $-\text{O}(\text{aryl})$ ,  $-\text{N}(\text{C}_0\text{-6alkyl})(\text{C}_0\text{-6alkyl})$ ,  $-\text{N}(\text{C}_0\text{-6alkyl})(\text{C}_3\text{-7cycloalkyl})$ ,  $-\text{N}(\text{C}_0\text{-6alkyl})(\text{aryl})$  substituents;

$\text{R}^4$  is  $-\text{C}_1\text{-6alkyl}$ ,  $-\text{C}_3\text{-7cycloalkyl}$ , heteroaryl, or aryl; optionally substituted with 1-5 independent halogen,  $-\text{CN}$ ,  $-\text{C}_1\text{-6alkyl}$ ,  $-\text{O}(\text{C}_0\text{-6alkyl})$ ,  $-\text{O}(\text{C}_3\text{-7cycloalkyl})$ ,  $-\text{O}(\text{aryl})$ ,  $-\text{N}(\text{C}_0\text{-6alkyl})(\text{C}_0\text{-6alkyl})$ ,  $-\text{N}(\text{C}_0\text{-6alkyl})(\text{C}_3\text{-7cycloalkyl})$ ,  $-\text{N}(\text{C}_0\text{-6alkyl})(\text{aryl})$  substituents;

A is  $-\text{C}_0\text{-4alkyl}$ ,  $-\text{C}_0\text{-2alkyl}-\text{SO}-\text{C}_0\text{-2alkyl}-$ ,  $-\text{C}_0\text{-2alkyl}-\text{SO}_2-\text{C}_0\text{-2alkyl}-$ ,  $-\text{C}_0\text{-2alkyl}-\text{CO}-\text{C}_0\text{-2alkyl}-$ ,  $-\text{C}_0\text{-2alkyl}-\text{NR}^9\text{CO}-\text{C}_0\text{-2alkyl}-$ ,  $-\text{C}_0\text{-2alkyl}-\text{NR}^9\text{SO}_2-\text{C}_0\text{-2alkyl}-$  or  $-\text{heteroC}_0\text{-4alkyl}$ ;

W is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>,  
 -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>,  
 5 -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or  
 -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents;

Y is optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>,  
 -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>,  
 -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>, -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>,  
 10 -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)R<sup>6</sup>, or  
 -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are combined to  
 form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>alkyl  
 substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl),  
 15 -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or  
 -N(C<sub>0-6</sub>alkyl)(aryl) groups;

R<sup>5</sup>, R<sup>6</sup>, and R<sup>7</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl,  
 heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>  
 20 alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

R<sup>8</sup> is -C<sub>1-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl, heteroaryl, or aryl; optionally  
 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>  
 cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl),  
 -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

25 B is -C<sub>0-4</sub>alkyl, -C<sub>0-2</sub>alkyl-SO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-SO<sub>2</sub>-C<sub>0-2</sub>  
 alkyl-, -C<sub>0-2</sub>alkyl-CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-NR<sup>10</sup>CO-C<sub>0-2</sub>alkyl-, -C<sub>0-2</sub>alkyl-  
 NR<sup>10</sup>SO<sub>2</sub>-C<sub>0-2</sub>alkyl- or -heteroC<sub>0-4</sub>alkyl;

R<sup>9</sup> and R<sup>10</sup> each independently is -C<sub>0-6</sub>alkyl, -C<sub>3-7</sub>cycloalkyl,  
 heteroaryl, or aryl; any of which is optionally substituted with 1-5 independent  
 30 halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>  
 alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), -N(C<sub>0-6</sub>alkyl)(aryl) substituents;

Z is -C<sub>3-7</sub>cycloalkyl, -heteroC<sub>3-7</sub>cycloalkyl, -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>  
 alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-</sub>





4. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein

Y is 2-pyridyl optionally substituted with 1-4 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>,  
 5 -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>,  
 -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>,  
 -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are  
 combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>  
 10 alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl),  
 -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or  
 -N(C<sub>0-6</sub>alkyl)(aryl) groups..

15 5. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein

X is phenyl optionally substituted with 1-5 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>,  
 -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>,  
 20 -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or  
 -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are combined to  
 form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1-6</sub>alkyl  
 substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl),  
 -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or  
 25 -N(C<sub>0-6</sub>alkyl)(aryl) groups.

6. The compound according to Claim 5, or a pharmaceutically acceptable salt thereof, wherein

Y is 2-pyridyl optionally substituted with 1-4 independent halogen,  
 30 -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>,  
 -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>,  
 -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>,  
 -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are  
 combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>  
 35 alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further

substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or -N(C<sub>0-6</sub>alkyl)(aryl) groups.

- 5                    7. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein

Y is 1,3-thiazolyl optionally substituted with 1-2 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>,  
 10 -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>, -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or  
 15 -N(C<sub>0-6</sub>alkyl)(aryl) groups.

8. The compound according to Claim 7, or a pharmaceutically acceptable salt thereof, wherein
- 20 X is phenyl optionally substituted with 1-5 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1-6</sub>alkyl  
 25 substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or -N(C<sub>0-6</sub>alkyl)(aryl) groups..

30

9. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein

W is -C<sub>0-6</sub>alkylaryl, or -C<sub>0-6</sub>alkylheteroaryl optionally substituted with 1-7 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl,

-OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>,  
 -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>, -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>,  
 -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents.

5                    10. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein

Y is pyrazolyl optionally substituted with 1-3 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>,  
 -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>,  
 10 -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>,  
 -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or  
 15 -N(C<sub>0-6</sub>alkyl)(aryl) groups.

11. The compound according to Claim 10, or a pharmaceutically acceptable salt thereof, wherein

20 X is 1phenyl optionally substituted with 1-5 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>,  
 -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>,  
 -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>,  
 25 -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1-6</sub>alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further substituted with 1-5 independent halogen, -CN, -C<sub>1-6</sub>alkyl, -O(C<sub>0-6</sub>alkyl), -O(C<sub>3-7</sub>cycloalkyl), -O(aryl), -N(C<sub>0-6</sub>alkyl)(C<sub>0-6</sub>alkyl), -N(C<sub>0-6</sub>alkyl)(C<sub>3-7</sub>cycloalkyl), or  
 -N(C<sub>0-6</sub>alkyl)(aryl) groups.

30

12. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein

Y is imidazolyl optionally substituted with 1-3 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkenyl, -C<sub>1-6</sub>alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>,

-C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>,  
 -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>,  
 -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are  
 combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1</sub>-  
 6alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 substituted with 1-5 independent halogen, -CN, -C<sub>1</sub>-6alkyl, -O(C<sub>0</sub>-6alkyl), -O(C<sub>3</sub>-  
 7cycloalkyl), -O(aryl), -N(C<sub>0</sub>-6alkyl)(C<sub>0</sub>-6alkyl), -N(C<sub>0</sub>-6alkyl)(C<sub>3</sub>-7cycloalkyl), or  
 -N(C<sub>0</sub>-6alkyl)(aryl) groups.

10 13. The compound according to Claim 12, or a pharmaceutically acceptable salt thereof, wherein

X is phenyl optionally substituted with 1-5 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1</sub>-6alkyl, -C<sub>1</sub>-6alkenyl, -C<sub>1</sub>-6alkynyl, -OR<sup>1</sup>, -NR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>,  
 -N(=NR<sup>1</sup>)NR<sup>2</sup>R<sup>3</sup>, -NR<sup>1</sup>COR<sup>2</sup>, -NR<sup>1</sup>CO<sub>2</sub>R<sup>2</sup>, -NR<sup>1</sup>SO<sub>2</sub>R<sup>4</sup>, -NR<sup>1</sup>CONR<sup>2</sup>R<sup>3</sup>, -SR<sup>4</sup>,  
 15 -SOR<sup>4</sup>, -SO<sub>2</sub>R<sup>4</sup>, -SO<sub>2</sub>NR<sup>1</sup>R<sup>2</sup>, -COR<sup>1</sup>, -CO<sub>2</sub>R<sup>1</sup>, -CONR<sup>1</sup>R<sup>2</sup>, -C(=NR<sup>1</sup>)R<sup>2</sup>, or  
 -C(=NOR<sup>1</sup>)R<sup>2</sup> substituents, wherein optionally two substituents are combined to  
 form a cycloalkyl or heterocycloalkyl ring fused to X; wherein the -C<sub>1</sub>-6alkyl  
 substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 substituted with 1-5 independent halogen, -CN, -C<sub>1</sub>-6alkyl, -O(C<sub>0</sub>-6alkyl), -O(C<sub>3</sub>-  
 20 7cycloalkyl), -O(aryl), -N(C<sub>0</sub>-6alkyl)(C<sub>0</sub>-6alkyl), -N(C<sub>0</sub>-6alkyl)(C<sub>3</sub>-7cycloalkyl), or  
 -N(C<sub>0</sub>-6alkyl)(aryl) groups.

14. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein


25 X is 3-pyridyl optionally substituted with 1-4 independent halogen, -CN, NO<sub>2</sub>, -C<sub>1</sub>-6alkyl, -C<sub>1</sub>-6alkenyl, -C<sub>1</sub>-6alkynyl, -OR<sup>5</sup>, -NR<sup>5</sup>R<sup>6</sup>,  
 -C(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -N(=NR<sup>5</sup>)NR<sup>6</sup>R<sup>7</sup>, -NR<sup>5</sup>COR<sup>6</sup>, -NR<sup>5</sup>CO<sub>2</sub>R<sup>6</sup>, -NR<sup>5</sup>SO<sub>2</sub>R<sup>8</sup>,  
 -NR<sup>5</sup>CONR<sup>6</sup>R<sup>7</sup>, -SR<sup>8</sup>, -SOR<sup>8</sup>, -SO<sub>2</sub>R<sup>8</sup>, -SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, -COR<sup>5</sup>, -CO<sub>2</sub>R<sup>5</sup>, -CONR<sup>5</sup>R<sup>6</sup>,  
 -C(=NR<sup>5</sup>)R<sup>6</sup>, or -C(=NOR<sup>5</sup>)R<sup>6</sup> substituents, wherein optionally two substituents are  
 combined to form a cycloalkyl or heterocycloalkyl ring fused to Y; wherein the -C<sub>1</sub>-  
 30 6alkyl substituent, cycloalkyl ring, or heterocycloalkyl ring each optionally is further  
 substituted with 1-5 independent halogen, -CN, -C<sub>1</sub>-6alkyl, -O(C<sub>0</sub>-6alkyl), -O(C<sub>3</sub>-  
 7cycloalkyl), -O(aryl), -N(C<sub>0</sub>-6alkyl)(C<sub>0</sub>-6alkyl), -N(C<sub>0</sub>-6alkyl)(C<sub>3</sub>-7cycloalkyl), or  
 -N(C<sub>0</sub>-6alkyl)(aryl) groups.


15. The compound according to Claim 1, consisting of:


- 5 1-methyl-3-[3-(5-pyridin-2-yl-2*H*-tetrazol-2-yl)phenyl]imidazolidin-2-one;  
 2-[2-(4-pyridin-2-ylphenyl)-2*H*-tetrazol-5-yl]pyridine;  
 2-[2-(4-pyridin-4-ylphenyl)-2*H*-tetrazol-5-yl]pyridine;  
 2-{2-[3-(1*H*-imidazol-1-yl)phenyl]-2*H*-tetrazol-5-yl}pyridine;  
 2-[2-(2-pyrazin-3-ylphenyl)-2*H*-tetrazol-5-yl]pyridine;  
 10 2-[2-(4-morpholin-3-ylphenyl)-2*H*-tetrazol-5-yl]pyridine;  
 2-{2-[3-(2*H*-tetrazol-5-yl)phenyl]-2*H*-tetrazol-5-yl}pyridine;  
 2-pyridin-2-yl-5-(5-pyridin-2-yl-2*H*-tetrazol-2-yl)benzonitrile;  
 or a pharmaceutically acceptable salt thereof.

15


16. The compound according to Claim 1, consisting of:








































































































or a pharmaceutically acceptable salt thereof.

17. A pharmaceutical composition comprising:  
 5 a therapeutically effective amount of the compound according to claim 1, or a pharmaceutically acceptable salt thereof; and  
 a pharmaceutically acceptable carrier.

18. The pharmaceutical composition according to claim 14, further  
 10 comprising i) an opiate agonist, ii) an opiate antagonist, iii) a calcium channel antagonist, iv) a 5HT receptor agonist, v) a 5HT receptor antagonist, vi) a sodium channel antagonist, vii) an NMDA receptor agonist, viii) an NMDA receptor antagonist, ix) a COX-2 selective inhibitor, x) an NK1 antagonist, xi) a non-steroidal anti-inflammatory drug, xii) a GABA-A receptor modulator, xiii) a dopamine agonist,  
 15 xiv) a dopamine antagonist, xv) a selective serotonin reuptake inhibitor, xvi) a tricyclic antidepressant drug, xvii) a norepinephrine modulator, xviii) L-DOPA, xix) buspirone, xx) a lithium salt, xxi) valproate, xxii) neurontin, xxiii) olanzapine, xxiv) a nicotinic agonist, xxv) a nicotinic antagonist, xxvi) a muscarinic agonist, xxvii) a muscarinic antagonist, xxviii) a selective serotonin and norepinephrine reuptake  
 20 inhibitor (SSNRI), xxix) a heroin substituting drug, xxx) disulfiram, or xxxi) acamprosate.

19. The pharmaceutical composition according to claim 18,  
 wherein said heroin substituting drug is methadone, levo-alpha-acetylmethadol,  
 25 buprenorphine or naltrexone.

20. A method of treatment or prevention of pain comprising the step of administering a therapeutically effective amount, or a prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

5

21. A method of treatment or prevention of a pain disorder wherein said pain disorder is acute pain, persistent pain, chronic pain, inflammatory pain, or neuropathic pain, comprising the step of administering a therapeutically effective amount, or a prophylactically effective amount, of the compound according to claim 1  
10 or a pharmaceutically acceptable salt thereof.

22. A method of treatment or prevention of anxiety, depression, bipolar disorder, psychosis, drug withdrawal, tobacco withdrawal, memory loss, cognitive impairment, dementia, Alzheimer's disease, schizophrenia or panic comprising the  
15 step of administering a therapeutically effective amount, or a prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

23. A method of treatment or prevention of disorders of  
20 extrapyramidal motor function comprising the step of administering a therapeutically effective amount, or a prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

24. The method of claim 23 wherein said disorder of extrapyramidal  
25 motor function is Parkinson's disease, progressive supramuscular palsy, Huntington's disease, Gilles de la Tourette syndrome, or tardive dyskinesia.

25. A method of treatment or prevention of anxiety disorders comprising the step of administering a therapeutically effective amount, or a  
30 prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

26. The method of claim 25 wherein said anxiety disorder is panic attack, agoraphobia or specific phobias, obsessive-compulsive disorders, post-

traumatic stress disorder, acute stress disorder, generalized anxiety disorder, eating disorder, substance-induced anxiety disorder, or nonspecified anxiety disorder.

27. A method of treatment or prevention of neuropathic pain  
5 comprising the step of administering a therapeutically effective amount, or a prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

28. A method of treatment or prevention of Parkinson's Disease  
10 comprising the step of administering a therapeutically effective amount, or a prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

29. A method of treatment or prevention of depression comprising the  
15 step of administering a therapeutically effective amount, or a prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

30. A method of treatment or prevention of epilepsy comprising the  
20 step of administering a therapeutically effective amount, or a prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

31. A method of treatment or prevention of inflammatory pain  
25 comprising the step of administering a therapeutically effective amount, or a prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

32. A method of treatment or prevention of cognitive dysfunction  
30 comprising the step of administering a therapeutically effective amount, or a prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

33. A method of treatment or prevention of drug addiction, drug abuse  
35 and drug withdrawal comprising the step of administering a therapeutically effective



amount, or a prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

34. A method of treatment or prevention of bipolar disorders  
5 comprising the step of administering a therapeutically effective amount, or a prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

35. A method of treatment or prevention of circadian rhythm and sleep  
10 disorders comprising the step of administering a therapeutically effective amount, or a prophylactically effective amount, of the compound according to claim 1 or a pharmaceutically acceptable salt thereof.

36. The method of Claim 35 wherein the circadian rhythm and sleep  
15 disorders are shift-work induced sleep disorder or jet-lag.

37. A method of treatment or prevention of obesity comprising the  
step of administering a therapeutically effective amount, or a prophylactically  
effective amount, of the compound according to claim 1 or a pharmaceutically  
20 acceptable salt thereof.

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/07074

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/4439; C07D 401/12

US CL : 546/268.4; 514/340

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 546/268.4; 514/340

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN CAS Online: CAPLUS, REGISTRY; EAST; WEST

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CAPLUS Accession No: 1986:511139, Kitaeva, et al. "Radioprotective and antitumor activity of some tetrazole derivatives," Khimiko-Farmatsevticheski Zhurnal (1986), Vol. 20, No. 5, pages 559-63.	1-17.

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier application or patent published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 23 June 2003 (23.06.2003)	Date of mailing of the international search report <b>17 JUL 2003</b>
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US Commissioner for Patents P.O. Box 1450 Alexandria, Virginia 22313-1450 Facsimile No. (703)305-3230	Authorized officer Alan Rouman <i>Alan L. Rouman</i> Telephone No. 703.308.1235